

**COMOX VALLEY REGIONAL DISTRICT  
SARATOGA BEACH AND BLACK CREEK AREAS  
(CVRD Electoral Area C)**

**INTEGRATED PEST MANAGEMENT PLAN  
PMP # 895-0002-26-31**

**MOSQUITO POPULATION MANAGEMENT AND CONTROL  
2026 - 2031**



Plan Prepared by  
[Duka Environmental Services Ltd.](#)

Surrey, BC

for

The Comox Valley Regional District  
Community Services  
Courtney, British Columbia

07 October 2025

# TABLE OF CONTENTS

<b>1.0</b>	<b>PEST MANAGEMENT PLAN SUMMARY .....</b>	<b>1</b>
1.1	Geographic Boundaries of this Pest Management Plan .....	2
1.2	Mosquito Biology.....	3
1.3	Need for Mosquito Control .....	3
1.4	Term of the Pest Management Plan (PMP).....	5
<b>2.0</b>	<b>MOSQUITO CONTROL PROGRAM BACKGROUND .....</b>	<b>5</b>
2.1	Primary Land Use.....	6
2.2	Mosquito Species Identified within the CVRD.....	6
2.3	Mosquito Control Program Implementation .....	8
2.4	Control Products (Larvicides) Proposed for Use.....	9
<b>3.0</b>	<b>CONTROL PROGRAM METHODOLOGIES .....</b>	<b>9</b>
3.1	Public Information and Education .....	10
3.2	Protection of Archaeological Sites.....	12
3.3	Mosquito Control Program Data Collection and Reporting.....	13
3.4	Surveying and Monitoring of Mosquito Populations .....	14
3.4.1	<i>Larval Mosquito Monitoring</i> .....	15
3.4.2	<i>Adult Mosquito Monitoring</i> .....	16
3.4.3	<i>Mosquito Development in Saratoga Beach and Black Creek</i> .....	17
3.5	Mosquito Control Options.....	19
3.5.1	<i>Physical Source Reduction and Site Modification</i> .....	20
3.5.2	<i>Biological Control</i> .....	21
3.5.3	<i>Biorational Control</i> .....	22
3.5.4	<i>Chemical Control</i> .....	24
3.6	Mosquito Control Program Operations.....	25
3.6.1	<i>Public, Worker, and Environmental Safety During Mosquito Control</i> .....	26
3.6.2	<i>Larval Mosquito Treatment Thresholds and Application Rates</i> .....	27
3.6.3	<i>Post Application Monitoring</i> .....	30
<b>4.0</b>	<b>QUALIFICATIONS OF PROGRAM PERSONNEL.....</b>	<b>31</b>
<b>5.0</b>	<b>LARVICIDE HANDLING AND APPLICATION .....</b>	<b>31</b>
5.1	Larvicide Transportation.....	32
5.2	Larvicide Storage.....	33
5.3	Larvicide Mixing, Loading and Application .....	33
5.4	Equipment Maintenance and Calibration .....	34
<b>6.0</b>	<b>CONTROL PROGRAM SYNOPSIS.....</b>	<b>34</b>
<b>7.0</b>	<b>REFERENCES AND LITERATURE CITED .....</b>	<b>35</b>
	<b>Figure 1: Saratoga/Miracle Beach; Larval Mosquito Development Site Locations.....</b>	<b>Appendix A</b>
	<b>VectoBac Product Information .....</b>	<b>Appendix B</b>
	<b>VectoLex Product Information .....</b>	<b>Appendix C</b>

## 1.0 PEST MANAGEMENT PLAN SUMMARY

The Saratoga Beach / Black Creek area of the Comox Valley Regional District (CVRD), Electoral Area C, and surrounding area has significant recreational and environmental value, providing residents and visitors with many outdoor summer activities and employment. Walking, running, cycling, bird watching, outdoor sports, golfing and gardening are just a few of these. Adult mosquito annoyance can often conflict with these activities and potentially impact public health. Besides the negative impacts on the lifestyle and well-being of residents, there can also be considerable economic impact from mosquito annoyance on local businesses. An integrated pest management (IPM) approach to mosquito population management and control can reduce overall annoyance levels and co-exist with these valuable resources.

This approach focuses on the timely detection and treatment of larval mosquito populations using biological products and methodologies. Where possible, and appropriate, physical or cultural controls are recommended, and implemented, that would reduce larval habitat and enhance or conserve natural mosquito predators. Where required, larval mosquito populations would be controlled using the biorational larvicide products VectoBac® 200G (*Bacillus thuringiensis var. israelensis*, Serotype H-14, Strain AM 65-52, PCP #18158) and VectoLex (*Lysinibacillus sphaericus*, PCP # 28008, 28009). A similar *Bti* product, AquaBac® 200G (*Bacillus thuringiensis var. israelensis*, Serotype H-14, Strain BMP-144, PCP #26863) may be used. All treatments would be completed in accordance with the methodologies and procedures prescribed in the BC Ministry of Environment-accepted Pest Management Plan (PMP) for Nuisance Mosquito Control, prepared by *Duka Environmental Services Ltd.* on behalf of the Comox Valley Regional District for the years 2026-2031.

Mosquito control services would be provided to residential and rural property owners, businesses, municipal and regional parks, sports fields, golf courses and other outdoor recreational and tourist facilities located within the boundaries of the CVRD Electoral Area C, and specifically the area described as Saratoga Beach and Black Creek. The goal of the annual mosquito control program is to reduce the potential for widespread adult mosquito annoyance for the benefit of residents, workers and visitors to the area.

The PMP is 'owned' by the Comox Valley Regional District. It would remain in place for the purposes of larval mosquito population management and control for a five year period, 15 April 2026 to 14 April 2031. The mosquito control program and methodologies developed within this PMP are a hybrid of approaches adapted from collaboration with mosquito and vector control professionals worldwide. It has been carefully and specifically designed for the unique conditions of the program areas and is a model of environmental compatibility. This PMP meets all the requirements of the Integrated Pest Management Act and will replace the previously approved, and soon to expire (April 2026) PMP # 895-0001-21/26. The mosquito control program proposed for the years 2026-2031 is largely unchanged from previous years and focuses on larval control and reduction of mosquito populations.


This PMP reviews mosquito biology, the types of larval mosquito habitats affecting the program area and the local mosquito species complex. An IPM approach to mosquito population management and control can reduce overall adult mosquito annoyance levels through education, prevention and biological controls. This PMP outlines the procedures and methodologies of an IPM approach which will reduce local mosquito populations for the purpose of reducing mosquito annoyance for area residents, workers and visitors.

The CVRD Mosquito Population Management and Control Program Pest Management Plan, PMP # 895-0002-26-31 (this document), is presented in a format which adheres to the requirements of British Columbia’s Integrated Pest Management Act and Regulation, including amendments, the Mosquito Management Sector Review Paper and BC Ministry of Environment, Indigenous Engagement: A Guide for Integrated Pest Management Act Proponents (2023). Copies of these documents may be accessed through the BC Ministry of Environment home page at [www.gov.bc.ca/gov/content/environment/pesticides-pest-management](http://www.gov.bc.ca/gov/content/environment/pesticides-pest-management). Common themes of larval development prevention and control necessary to achieve program goals, while ensuring environmental conservation, are repeated throughout this document.

A professional, experienced, environmental services firm (the consultant) would be retained by the Comox Valley Regional District to deliver these very specialized services and to ensure adherence to the Pest Management Plan. The consultants for the CVRD annual mosquito control program would have Registered Professional Biologists (R.P.Bios.) as program managers and senior biologists. All program personnel would be appropriately certified as pesticide applicators with the BC Ministry of Environment, Integrated Pest Management Program.

Public relations and ongoing program education would be accomplished through regular contacts with residents, businesses and community visitors. Information on mosquitos, their control, and prevention, is available to the general public in a variety of forms including informational brochures, websites, newspaper articles, open-houses, council meetings and farmer’s markets etc. Resident requests for service are followed up with telephone contact and site inspection. Physical reduction, elimination or alteration of larval mosquito development habitats is an important aspect of the control program. Wherever possible, and practical, property owners are advised of measures they can complete to reduce mosquito development and adult mosquito annoyance.

### 1.1 Geographic Boundaries of this Pest Management Plan




The Saratoga Beach and Black Creek area is a rural residential community located midway between the City of Courtney and City of Campbell River, approximately 30 minutes from each community by car. Located within the CVRD Electoral Area “C”, this seaside community borders the Strait of Georgia to the east and the Oyster River along the northern boundary. Highway 19A (old Island Highway) is the western-most boundary of the mosquito abatement/control service area.

The Saratoga Beach and Black Creek area has significant recreational and environmental value, providing residents and visitors with many outdoor summer activities including hiking, walking, camping, fishing, boating, cycling, golfing, gardening, etc. Saratoga Beach offers one of the sandiest beaches in the area and has several parks, both Regional and Provincial, several campgrounds and resorts, Bed and Breakfasts and the Saratoga Beach Golf Course. A map indicating the boundary of the proposed PMP area and current identified larval mosquito development sites can be found in Appendix A at the end of this document.

## 1.2 Mosquito Biology

Mosquitos are found world-wide in standing water of all possible descriptions. Mosquitos belong to the order Diptera, along with other pests such as the common house fly and the black fly. There are over sixty species common to Canada and over thirty are found in British Columbia. Mosquitos undergo four distinct development stages: egg, larvae, pupae and adult. Larvae and pupae are aquatic. Eggs are laid on the water surface or on soil and vegetation adjacent to water. The eggs of some species of mosquitos, such as *Aedes*, can survive for upwards of 20 years and will hatch after a period of winter freezing and upon being inundated with water.



Mosquito larvae undergo four larval instars (or moults), each time emerging larger, but virtually unchanged from the previous instar. This is the feeding stage of the aquatic mosquito. The mosquito pupa, like a butterfly chrysalis, is a non-feeding stage and is where the once aquatic, larval mosquito undergoes metamorphosis to emerge as the winged, terrestrial adult mosquito. Adult mosquitos feed on plant nectar and other fluids, and it is only the female which requires a necessary blood meal to complete the development of her eggs. Any container, pond, or depression, either natural or manmade, which is capable of holding water for several days, to a few weeks, can provide development habitat for larval mosquitos.

Mosquito development occurs in a wide range of larval habitats ranging from coastal saltmarshes, snowmelt and precipitation-influenced flood and seepage water pools and channels along rivers and lakes, to freshwater ponds, marshes, ditches and catch basins. Birdbaths, plugged rain gutters, livestock watering troughs and any man-made container capable of holding water for a period of 7 to 21 days can provide suitable larval mosquito habitat.

## 1.3 Need for Mosquito Control

The purpose of the annual mosquito population management and control program is to provide residents, workers and visitors of the Saratoga Beach and Black Creek area of the CVRD Electoral Area C with relief from extreme and/or persistent adult mosquito annoyance using an Integrated Pest Management (IPM) approach to surveillance and population management. The control program is not intended to, nor is it possible to, eradicate local mosquito populations. An effective,

pro-active mosquito control program which focuses on the identification, prevention, or timely control of larval mosquito populations, also contributes to the protection of public health.

In addition to negative impacts on the lifestyle and general health of residents, a large population of mosquitos can have a negative economic impact on local businesses. Worker safety, comfort and efficiency can be compromised by adult mosquito annoyance and distraction. Reduced use and enjoyment of hotel and restaurant outdoor patios, sports fields, golf courses, campgrounds and cycling or hiking trails by residents and area visitors directly affects business operations and revenues.

Although not a common occurrence in most areas of British Columbia, mosquitos are capable of transmitting (vectoring) diseases. A well organized and effective larval mosquito control program is important to limit the potential for both, widespread adult mosquito annoyance, and potential for disease transmission. Despite the best of efforts though, some adult mosquito annoyance may still occur during the months of June through August, and residents are encouraged to avoid areas of mosquito harbourage (typically treed, forested or landscaped areas) during certain times of day, and to use repellents and approved adult mosquito control devices and products as per label directions.

Mosquitos are best known as vectors of 'tropical' diseases such as malaria and yellow fever. Although these exotic afflictions are extremely rare in British Columbia, mosquitos can still pose a serious health concern. Extreme allergic reactions or secondary infections from mosquito bites can occasionally require hospitalization. Diseases such as canine heartworm, Western Equine Encephalitis (WEE) and West Nile virus (WNV) are transmitted from some mosquito species to family pets, humans, and livestock. A few years ago (2019), the mosquito-associated flavivirus disease caused by Zika virus (ZIKV) became a prominent health concern in several areas of the world, including the southern USA. Locally in British Columbia, the recent discovery (July 2025) of a "cluster" of Jamestown Canyon virus and Snowshoe Hare virus infections in several residents of Squamish, BC (BCCDC, July 2025) has increased concerns for invasive species and disease range expansions.

Since mosquitos capable of vectoring diseases to man are often the source of annoyance (human-biting), the control of mosquito populations known to cause nuisance also contributes to the protection of public health by controlling mosquito species also having the potential to vector disease. A few years ago, the mosquito-associated flavivirus disease caused by Zika virus (ZIKV) became a prominent health concern in several areas of the world, including the southern USA. The current status of WNV and ZIKV in British Columbia, Canada and elsewhere in North America is available at [www.BCCDC.ca](http://www.BCCDC.ca) and Health Canada at [www.canada.ca/en/health-canada](http://www.canada.ca/en/health-canada) or [www.Hc-sc.gc.ca](http://www.Hc-sc.gc.ca).

The BC Centre for Disease Control (Vancouver) and local health authorities are responsible to coordinate the ongoing surveillance, identification and reporting of WNV, ZIKV, and new or emerging diseases, and their mosquito vectors. As part of this planning the BCCDC has developed the Arbovirus Surveillance and Response Guidelines for British Columbia, and the BCCDC has a provincial database containing all mosquito, bird and human health surveillance data relating to WNV with ongoing updates for other, new vector mosquito species.

Due to the low and stable incidence of WNV it was decided by the BCCDC in the fall of 2014 that it was no longer necessary to conduct active surveillance of mosquitos or other indicators. The provincial decision to eliminate this surveillance was reached at the BC Communicable Disease Policy Advisory Committee meeting in February 2015. Human clinical testing will continue. Human clinical testing continues as part of routine blood donor collection programs. Dead birds such as crows and ravens are tested when WNV is suspected and horses are also tested, on a case-by-case basis, by provincial veterinarians.

#### **1.4 Term of the Pest Management Plan (PMP)**

A five-year period, extending from 15 April 2026 to 14 April 2031.

The designated contact for this Plan is Mr. Mark Harrison, Manager of Parks, Comox Valley Regional District, 770 Harmston Avenue, Courtney, BC V9N 0G8. Telephone #250-334-6067 or Email [mharrison@comoxvalleyrd.ca](mailto:mharrison@comoxvalleyrd.ca).

## **2.0 MOSQUITO CONTROL PROGRAM BACKGROUND**

In response to several seasons of resident, business and facility operators reports of extreme adult mosquito nuisance the CVRD commissioned two development site surveys. Completed in 2017 and 2020 they identified the predominant water courses and potential larval mosquito development habitats.

The largest source of mosquito development identified in the Saratoga Beach and Black Creek area is a salt water (tidal) influenced slough (Driftwood Marsh) extending from Black Creek northwards in an old channel/back dune area approximately 10-30m wide. With a length of some 900 - 1000m, the total area of this site is ~20,000m<sup>2</sup> (2.0 ha). A similar, smaller (< 2,000m<sup>2</sup>) salt water influenced habitat/marsh exists adjacent to the Pacific Playgrounds Resort and Marina at the end of Clarkson Avenue. Figures 1 and 2 present the primary sources of larval mosquito development (salt marshes) and distribution of adult mosquito populations sampled and identified as part of the 2020 Mosquito Development Site Survey.

Additional, potential larval mosquito development habitats includes local golf course ponds, roadside ditches and other freshwater water accumulations from seepage and precipitation water run-off in forested areas behind the Miracle Beach School and within the residential and business areas of the community. Ongoing and routine surveys to be conducted as part of the annual, operational mosquito population management and control program would identify other sources of larval mosquito development.

The mosquito control program and methodologies (the PMP) developed for the CVRD are a hybrid of approaches developed through collaboration with mosquito and vector control professionals worldwide. It has been carefully and specifically adapted for the unique conditions of the program area and is a model of environmental compatibility. A variety of monitoring and control methods, including physical site reduction or modification and the use of biological control products support

the principles of an Integrated Pest Management (IPM) approach to mosquito control. They are the most effective means of reducing adult mosquito populations and the potential for annoyance or disease transmission. This IPM protocol consists of five components:

- 1) Public Education. Explains mosquito biology, the program, and how the public can contribute to successful operations;
- 2) Surveillance, identification, and distribution of mosquito species;
- 3) Timely implementation of mosquito controls and preventative measures;
- 4) Adaptive management of operations during a season in response to observations; and
- 5) Program evaluation and assessment to ensure sustainable, effective controls have been achieved and the need for modification in future seasons.

The annual mosquito population management and control program focuses mosquito surveillance and control efforts in areas where larval populations are known to occur and where past, occasionally notable, adult mosquito annoyance was documented. Mosquito control services are provided to residential and rural property owners, businesses, municipal and regional parks, sports fields, campgrounds, golf courses and other outdoor recreational and tourist facilities.

## **2.1 Primary Land Use**

The primary land uses of the areas contained within the control program are agricultural (hobby farms), residential, light industrial and commercial properties (restaurants, garages, shopping plaza), recreational (golf courses, passive parks, sport and playing fields, etc.), Miracle Beach Provincial Park, Driftwood Marine Park and other forested and undeveloped lands. In addition to organized sports activities, outdoor summer recreational activities include walking, hiking, photography, golfing, camping, fishing, boating, sight-seeing and bike riding.

## **2.2 Mosquito Species Identified within the CVRD**

Mosquito development occurs in a wide range of open water larval habitats, ranging from snowmelt and precipitation-influenced flood and seepage water pools and channels along rivers, streams and lake shorelines, to permanent ponds, marshes, ditches and similar water-holding depressions. Bird baths, plugged rain gutters, livestock watering troughs, irrigation and surface water run-off collection ponds and any other man-made container or excavation capable of holding water for a period of at least 7 to 10 days can provide suitable larval mosquito habitat. Left undetected, larval mosquitos will complete their development to adult within this time span. Larval populations are sampled using a standard 350 ml dipper and area recorded as # larvae/350 ml dip sample, and by their age class, 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, 4<sup>th</sup> or pupae.

Eleven species of mosquito have been collected as either larvae or adults in the Saratoga Beach / Black Creek area. All of the following species present in the region, except *Culex territans* (which feeds on amphibians), can be a nuisance and many species, particularly those which bite birds and mammals other than man, are also capable of vectoring diseases, such as Western Equine Encephalitis and West Nile virus.

Mosquito pest species collected locally from the region include:

*Aedes cinereus*

*Aedes sticticus*

*Culex pipiens*

*Aedes dorsalis*

*Aedes vexans*

*Culex tarsalis*

*Aedes triseriatus*

*Anopheles punctipennis*

*Culex territans*

*Aedes provocans*

*Culiseta impatiens*



Larval Mosquitos in 350ml Dipper  
(200 larvae / dip, 2<sup>nd</sup> & 3<sup>rd</sup> instar)

The large majority of mosquito species collected from the area, over 90%, are a complex of *Aedes* mosquitos often containing large populations of *Aedes dorsalis*, the “Salt Marsh Mosquito”. All *Aedes* mosquitos are aggressive biting pests which prefer flooded, temporary or recurring habitats such as tidal salt marshes, alkaline pools, over-irrigated fields, ditches, marshes, snowmelt, river flood and seepage-filled pools in woodlands, sloughs and depressions in cleared or logged areas, golf courses and walking, cycling trails. *Aedes* mosquitos lay their eggs in moist soil along the edges of recently flooded areas where the eggs can lay dormant for many years. Following a period of wetting, and drying, eggs become “primed” to hatch. *Aedes* larvae, once inundated, can hatch out in large numbers with populations typically ranging from 50-100

larvae/dip sample, although + 200 larvae/dip sample isn’t uncommon. This is comparable to upwards of 2000-3000 mosquitos per square meter of surface water. *Aedes* mosquitos are typically the most numerous during the first half of the season, from late April through early July. Receding river levels, increasing ambient temperatures, evaporation and decreased precipitation causes many of these habitats to dry, drain and numbers upwards of 200 larvae/350ml dip disappear over the course of a typical summer.

*Culex* and *Culiseta* mosquitos comprise the remaining mosquito pest species for the area. They typically develop later in the season, from June through August, and require a different set of cues to initiate the onset of larval development. They prefer permanent and slow-draining, or frequently refilled sites including natural and man-made ponds, ditches and containers such as stored tires, boats and buckets or livestock watering troughs. Larval populations typically range from 1-20 larvae/dip sample and multiple, or recurring hatches are possible with additional egg laying by female adults. Although their populations and individual development sites are not usually as large as the synchronous hatching *Aedes* mosquitos, *Culex* and *Culiseta* can be a source of reportable annoyance since their preferred habitats are common to residential, commercial, recreational and agricultural properties. Species such as *Culex tarsalis* are able to withstand a high degree of pollution and can inhabit areas with high organic content, including septic field seepage, sewage lagoons and livestock hoof prints around barns, feed lots and along creeks. *Culex pipiens*, the “house mosquito”, can use a large variety of freshwater habitats including manmade containers and they are the predominant mosquito developing in catch basins.

All of the species listed above are able to develop as multiple hatches (generations) during the season and most of them are capable of causing reportable and often extreme annoyance, particularly *Aedes*. Locally collected *Aedes dorsalis* is also a potential West Nile virus (WNV) vector. *Culex* and *Culiseta* mosquitoes are not only a source of annoyance, but they too are also recognized as vectors of several diseases, including WNV. *Culex tarsalis*, *Culex pipiens* and *Culiseta incidens* *Culiseta inornata* are identified by the BC Centres for Disease Control (BCCDC) and the Centers for Disease Control (Atlanta, USA) as three of the primary vector species of WNV in North America. Control of locally occurring *Aedes*, *Culex* and *Culiseta* mosquitoes not only prevents widespread nuisance for the benefit of residents, businesses and visitors, but also contributes to the protection of public health. Mosquito surveillance, collection and identification are components of an ongoing operational control program. The mosquito species listing and development site database would be updated as required.

### 2.3 Mosquito Control Program Implementation

In response to resident, worker, and visitor reports of recurring adult mosquito annoyance, the Comox Valley Regional District has undertaken the development and delivery of an annual mosquito population management and control program for residents, workers and visitors, of the Saratoga Beach and Black Creek areas. The operational program described within this PMP will provide environmentally sound, and sustainable, mosquito control using an IPM approach. This methodology incorporates public education, development site identification and categorization, surveillance, prescriptions for site alteration or modification, and where required, larval mosquito control completed using the safest, most effective biological control agents available.

Ongoing mosquito development site surveys, monitoring and identification of larval and adult mosquito specimens updates the local mosquito species complex and development site database. Identified mosquito habitats are monitored throughout the season, typically from late April through August, to assess the abundance and species of mosquitoes developing in them. New Jersey or CDC (Atlanta) light traps and standardized mosquito biting and landing counts can be used to sample and monitor adult mosquito populations.

Two tidally influenced (salt water) development sites (Driftwood Marsh) were identified and confirmed as the primary sources of *Aedes dorsalis* and the reported adult mosquito nuisance, (Figure 1). Other potential habitats are present in the area as evidenced by adult mosquito sample collections from 2020 through 2025. In addition to *Aedes dorsalis*, ten species of mosquitoes, all of which use freshwater habitats, were collected as larvae or adult specimens from 2020 - 2025. Potential habitats include roadside ditches, irrigation or golf course display ponds, natural and man-made ponds on private and commercial properties. Salt marsh habitats which may contain a hundred or more individual, temporarily filled ponds, ditches and depressions. Flooding and seepage water accumulations from spring and summer tides provide extensive habitat for repeated *Aedes dorsalis* larval development in area salt marshes. Stagnant and non-flowing ponds and ditches, most of them manmade or influenced, provide freshwater larval mosquito development habitat and often have the greatest diversity of species.

Other habitats such as bird baths, buckets, stored boats, livestock watering troughs, tires are not treated as part of routine control program operations. When discovered, physical control of these habitats can be easily accomplished by removal of the container or for bird baths or watering troughs, regular drainage and refilling. This prevents larval mosquito development and subsequent adult mosquito annoyance. Public education activities encourage property owners to survey their properties and identify these types of habitats for removal or routine management.

Adult and larval mosquito population monitoring is conducted as part of ongoing operational mosquito management and control programs. This allows for an assessment of larval control effectiveness in reducing nuisance mosquito populations, updates the local species record and contributes to the larval mosquito development site database and information.

#### **2.4 Control Products (Larvicides) Proposed for Use**

The Comox Valley Regional District Mosquito Population Management and Control Program reduces adult mosquito populations by focusing on the identification and suppression of larval mosquito development using an IPM approach. This approach includes site modification or elimination, the conservation and enhancement of natural predators and controls, and when these are ineffective, or inefficient, the targeted use of biological and biorational control products.

The bacterial mosquito larvicide VectoBac 200G (PCP # 18158), *Bacillus thuringiensis var. israelensis* (*Bti*) Serotype H-14 has been routinely used in the annual program for 5 years. Other bacterial larvicide products, VectoLex CG (PCP # 28008) and VectoLex WSP (PCP # 28009), are made with *Lysinibacillus sphaericus*. A similar product, AquaBac 200G (PCP # 26863), made with different strains of the bacterium, *Bacillus thuringiensis var. israelensis* (*Bti*) Serotype H-14. VectoBac 200G and AquaBac 200G, produced by *Bacillus*, and VectoLex (CG & WSP) produced by produced by *Lysinibacillus*, all contain spores and crystals produced by the naturally occurring soil bacterium, and are classified as a biorational control products.

They are very specific to mosquito larvae and have no impact on non-pest insects, fish, amphibians, birds, reptiles and mammals. Extensive product information can be found for VectoBac and VectoLex at the manufacturer's website [www.valentbiosciences.com](http://www.valentbiosciences.com). Product information can also be found through the Health Canada, Pest Management Regulatory Agency (PRMA) website <https://www.canada.ca/en/health-canada/services/consumer-product-safety/pesticides-pest-management.html> and the Pesticide Label Search <https://pr-rp.hc-sc.gc.ca/lr-re/index-eng.php>. Section 3.4.3 Biorational Control, discusses the products VectoBac (*Bti*) and VectoLex (*Lsph*) further.

### **3.0 CONTROL PROGRAM METHODOLOGIES**

The objective of the annual mosquito control program is to reduce the potential of widespread adult mosquito annoyance for residents, workers, and visitors to CVRD Electoral Area C, and specifically the communities of Saratoga Beach and Black Creek . A program of this scope is not intended to eliminate the local mosquito population. The total eradication of a widespread, fecund insect pest is not feasible nor desirable. The potential impacts of control products and activities, combined with

a need to coexist with a delicate aquatic habitat, necessitates that an integrated approach to mosquito control be undertaken. This approach requires an assessment of the problem, an in-depth understanding of factors influencing the situation, followed by the use of appropriate control.

Control of or prevention of larval mosquito development is preferred over control of the often widely dispersed and mobile adult mosquito. Mosquito larvae are concentrated in one place, must remain there for 7-21 days, and are very susceptible to the biorational control (larvicide) products, VectoBac and VectoLex. Drainage, filling of depressions, restoration of flow in ditches or other physical alterations to appropriate larval mosquito development sites is the preferred and permanent control method.

Physical control can be integrated into local public works and construction activities such as roadside grading, ditch maintenance and cleaning. For home and business owners it can include the removal of water-holding containers such as buckets and unused pools, or the regular draining and refilling of livestock watering troughs and bird baths. Only the most environmentally compatible, least toxic and persistent control products would be deployed for use within this annual program. Specifically, the biorational larvicides, VectoBac 200G and VectoLex both made with the *Bacillus* sp. bacterium are the larval control products of choice.

Adult mosquito populations are monitored at select locations within the control program and in response to resident requests for service. When adult mosquito annoyance is identified, surveillance for potential, and unknown development sites can be undertaken, and controls completed, to reduce mosquito populations before they can disperse to cause increased annoyance.

Routine adult mosquito control applications (adulticiding) for the purposes of nuisance mosquito control **are not** a component of the CVRD Mosquito Population Management and Control Program and this Pest Management Plan. The operational components of the PMP and the successful, annual, Mosquito Population Management and Control Program may include the following activities, as detailed in Sections 3.1 through to Section 4.5.

### 3.1 Public Information and Education

The general public must be advised of control program efforts in their area and provided with the opportunity to have input to their mosquito control program. Public input is invaluable to any community function, and it is a key component of all successful, pro-active mosquito control programs. This is essential since, in the final assessment, it is the general public which must be satisfied with control efforts.

The *Integrated Pest Management Act and Regulation* requires public notification of Pest Management Plan preparation through newspaper notices which must be published twice in a two week period starting at least 45 days before submission of a notice confirming that a pest Management Plan has been prepared according to the legislation. The



general public, first nations and other stakeholders are invited through these advertisements to provide comments on the PMP and to consult with the PMP holder or his or her designate, on PMP contents and the proposed mosquito population management and control program. In addition, those individuals or groups which had requested information or who have supplied input when the local mosquito control program was last advertised and approved are contacted directly each time the PMP is renewed.

The annual mosquito population management and control program is well known and supported by area residents and businesses. Its highly visible nature using field biologists working along roadsides, fields, in parks, golf courses, along dykes and lake shorelines has ensured that property owners, facility managers and residents remain familiar with their annual program. In annual operation for five years, it has been providing mosquito surveillance, monitoring and larval control services for the benefits of residents, businesses and visitors to the area.

Considerable value can be obtained through exposure of the control program and interactions with the public. For example, public contact can result in the locating of new mosquito development sites thus augmenting efficacy. Residents are typically encouraged to contact control program consultants or CVRD offices (250-334-6000 or [www.comoxvalleyrd.ca](http://www.comoxvalleyrd.ca)) to report potential sources of larval mosquitos (a waterbody) or adult mosquito annoyance. Suggestions for physical removal or source reduction on private property allow the owner to participate on a smaller scale. Once accomplished, physical source reduction, especially the removal of artificial containers, grading of depressions or filling of tire ruts eliminates the need for further attention.

Movement of adult mosquitos, either by active flight or passively by wind, from outside of treated areas into built up and developed areas is always a possibility. Public education further encourages residents and businesses to undertake actions for excluding adult mosquitos and modification of personal behaviours which will reduce the potential for annoyance. Through eliminating development sites on their property and learning to reduce adult mosquito annoyance residents can actively participate in their program. In addition to providing residents with information on how they can reduce larval development and annoyance around their properties, education initiatives help residents understand that the control program can only suppress mosquito populations, not eradicate them, and that some adult mosquito annoyance may be anticipated at certain locations, times of day and during some years.

Examples of some various public education and information initiatives which could be employed throughout the CVRD in the following years include:

- Informational Brochures – these review mosquito biology and control, mosquito “myths”, program operations and contact information for program biologists.
- Cardboard Doorknob Hangers – These “Sorry we missed you” doorknob messages are left when residents aren’t home during property inspections. They summarize field technician observations and have return contact information for resident use.



- Laminated posters – durable. Can provide basic information on protection from annoyance. Installation along walking trails, picnic and camping areas is possible.
- Web-based program information and service contact details.
- Social media – Facebook, Twitter – another method of public access and information sharing.
- Newspaper Display Advertisements – placement in local newspapers from April – September. Provides public information relevant to each month and program contact/access information.
- Newsmedia interviews/articles – TV, Radio provides public with updates on program operations and status, mosquito biology and additional public outreach.
- Presentations at Council meetings.
- Information at Open houses, farmers markets.



As part of annual control program start-up in early April and May, program personnel contact property owners, residents and facility operators in the database to determine site status and confirm program participation and property access. Ongoing interactions and conversations with property owners, residents and the general public provides opportunities to discuss program operations and the distribution of public education and outreach materials and information.

Occasionally individuals may wish to be excluded from the mosquito control program for personal reasons. A record of "AVOID" areas is maintained and updated as required. Meetings and input with concerned residents and special interest groups ensures that activities of control personnel do not conflict with those of residents. By staying informed of community events such as baseball games, tournaments, rodeos and the like, control personnel can increase efforts prior to an event to reduce potential adult mosquito annoyance.

The cooperation and support of local businesses, farmers, business and facility operators and other property owners is indicative of true community spirit and support for a successful program which benefits workers, residents and visitors to the area. Prevention of adult mosquito annoyance through pro-active, larval mosquito control provides significant benefit to outdoor worker and recreational uses.

### 3.2 Protection of Archaeological Sites

Archaeological sites on both public and private land are protected under the Heritage Conservation Act (HCA) and must not be altered without a permit. Archaeological sites are non-renewable and have cultural, historical, scientific and educational value. The HCA automatically protects all archaeological sites that predate AD 1846, with exception of burial sites and rock art sites which are

protected regardless of age. Any individuals working in the annual Mosquito Surveillance and Control Program that believe they may have encountered materials or items of archaeological importance will follow the procedures below:

- All work in the vicinity of the items/objects will cease immediately and any archaeological and/or human remains will not be disturbed.
- Will contact their supervisor/program manager.
- No excavation or removal of soil from the area will occur.
- Will isolate, mark, and protect the area from disturbance.
- Take pictures of the artifact, the immediate and adjacent areas.
- Note location (GPS coordinates, location description) and leave all discoveries in place.
- The CVRD and provincial Archaeology Branch (email: Archaeology@gov.bc.ca, or 250-953-3334) will be contacted.

### **3.3 Mosquito Control Program Data Collection and Reporting**

The environmental consultant (contractor) managing the annual mosquito population management and control program for the CVRD is responsible to follow the data collection and reporting requirements of the PMP and the *Integrated Pest Management Act and Regulations*.

The CVRD Community Services office would be regularly informed of control program activities of this contractor/consultant through personal contact, telephone, facsimile or e-mail with program managers and field personnel. In addition, written progress reports summarizing weather conditions, surveying and monitoring results, treatment areas and interactions with the public are typically prepared by the program consultants and submitted to the District on regular basis during operational phases of the control program.

At the conclusion of each annual mosquito control program season a summary report detailing all activities and pesticide treatments completed under the PMP is provided by the consultant to the CVRD. All pesticide use reporting required under the Integrated Pest Management Act, the approved PMP and as requested during the season by government regulatory agencies, including the BC Ministry of Environment would be completed by the consultant (as agents for the CVRD), as requested and necessary.

At a minimum, the contractor would maintain the following information for their use in managing the program and to complete the reporting and information requirements of the CVRD, the PMP and the Pesticide Management Program, the IPM Act and Regulations, and the BC Ministry of Environment:

- A mosquito development site database with information including property ownership, address, contact telephone number, public access information (paths, trails, roadways), development site maps, photographs, GPS identification, records of monitoring and treatment activities, pesticide use daily operation records and other relevant information related to the control program.

- A record of properties identified as 'AVOID' areas, where the owner or residents have indicated through telephone, written, verbal (in person conversation) or electronic (e-mail, facsimile) communication with the CVRD or its consultants, their wish to be excluded from the mosquito control program.
- A list and/or maps identifying 'AVOID' areas of environmental sensitivity, including provincial or regional parks, habitat conservation areas and other identified or designated speciality management areas.
- When the status of a waterbody or other area of potential environmental concern (e.g. bird nesting sites) is unknown, a local representative of Fisheries and Oceans Canada (DFO) or the BC Ministry of Environment (BCMOE), or other agencies where appropriate, may be consulted.

The development site database and avoid area lists are updated during each field season when control program personnel meet with residents, owners and operators of the farms, businesses and recreational facilities. Property ownership, access, development site status, areas of concern and control program operations are reviewed at this time. Regular contact is maintained with these individuals throughout the season to provide updates on control program operations and opportunities for input and comment on the control program. Ongoing activities related to surveying, monitoring and mosquito control operations are recorded in the historical data section of the database as they occur. Database information would be used to respond to any requests for program information from the public, District representatives or government regulatory agencies.

The consultant is responsible to ensure that all private and personally identifiable information within the database is managed in a secure, ethical, and legally compliant manner. This includes, but is not limited to, names and contact information linked to individuals or private properties, and any other data that could reasonably be used to identify a person. All handling of such information must adhere to the requirements set out in both the Personal Information Protection and Electronic Documents Act (PIPEDA) and the Personal Information Protection Act (PIPA). These Acts govern how private-sector organizations collect, use, disclose, and protect personal information in the course of commercial activities.

### **3.4 Surveying and Monitoring of Mosquito Populations**

As part of the annual program start-up, and throughout the season, program field biologists conduct regular, comprehensive surveys of the program area, as appropriate. The goal of these surveys is to confirm the extent and locations of existing, known mosquito development sites and to identify any new, or previously undetected, larval habitats. Surveying and monitoring of larval development sites (always waterbodies) determines the presence of larval mosquitos, the need for control and allows for regular update of the database. Where observed, larvae are collected and enumerated using a standard 350 ml white larval mosquito dipper. Preserved larval specimens are identified to the species when possible.

Mosquito development varies from year to year and throughout the season depending on environmental conditions and habitat availability. Environmental cues interact to affect both the timing and magnitude of mosquito development, and adult mosquito survival. Provincial and regional snowpack accumulations, river levels, tidal heights, precipitation and temperatures are reviewed as necessary to ensure timely surveying to detect mosquito development.

Monitoring and correlation of fluctuating temperatures and precipitation levels over several seasons allows for the determination of 'thresholds' which aid in the prediction of larval development and distributions. Review of winter snowpack accumulations, summer weather conditions, and tidal levels combined with a sound knowledge of mosquito biology and local development site types is necessary to ensure surveying and monitoring activities occur to detect mosquito development. Failure to timely survey and monitor could allow unchecked development of larvae which will result in adult mosquito annoyance. Larval habitats would be monitored throughout the season to assess the relative abundance and species of larval mosquitos found in these habitats. When investigating reports of adult mosquito annoyance or potential larval development sites, a thorough survey of each area would be performed to locate the source of annoyance, and any previously unidentified larval habitat.

### **3.4.1 Larval Mosquito Monitoring**

Surveying and monitoring of larval development sites (always waterbodies) determines the presence of larval mosquitos and the need for control. Larval habitats would be monitored throughout the season using a standard 350ml white larval mosquito dipper to assess the relative abundance and species of larval mosquitos found in these habitats. Routine sampling of development habitats is ideally completed on a 6-10 day basis, depending on conditions and observations, throughout the operational season, typically mid-April to late August.

Larval mosquito populations as small as one larva per 350ml dip sample in an area as small as a backyard swimming pool (5m x 10m) can produce thousands of adult mosquitos over the course of a season. Located adjacent to established outdoor recreational facilities including golf courses, sports parks, picnic areas, campgrounds and nearby residential and commercial areas, seepage and snowmelt influenced sites, including river level-influenced sites, are a major potential source of mosquito annoyance, and a primary focus of the annual mosquito control program

Pre-treatment surveys determine the extent of larval development which ensures that control measures are directed only to those areas containing larvae. In addition to providing pre-application information essential to timely control applications, surveying and monitoring following treatment, 'post-treatment monitoring' allows for an evaluation of the degree of control achieved from a particular application. Environmental compatibility and cost effectiveness of a control program is dependent on proper pesticide use through the application of control measures directed only to those areas requiring them. Post-treatment monitoring to confirm larval mortalities is typically completed within 2-96 hours of larvicide (VectoBac 200G and VectoLex CG) application.

### 3.4.2 Adult Mosquito Monitoring

To objectively measure the success and effectiveness of larviciding efforts in reducing adult mosquito populations, two internationally accepted sampling methods are employed. The first, a standard biting/landing count, measures the number of mosquitos which land, to bite, on the exposed forearm (from wrist to elbow) in a one minute period. Adult biting counts of three or more per minute, measured between the wrist and exposed forearm, is intolerable for most people. Beyond three bites per minute, outdoor enjoyment and worker performance and safety are affected, and negative economic impacts on recreation and tourism can be expected.

Although it is the accepted world-wide standard, it must be noted that bite counts are not without bias. Clothing and body physiology make some people more or less attractive than others. Also, daily timing for collection is crucial as mosquitos are most active at dusk and dawn, when temperatures are lower and humidity generally higher. For these reasons, collection timing, locations and clothing worn by the observer are standardized as much as possible. When reviewed in conjunction with anecdotal reports from residents, this data is a useful measure of mosquito annoyance levels and facilitates the collection of mosquito species that actively seek a human blood meal.




*CDC Light Trap*

The second method used for adult mosquito population assessments uses either Standard New Jersey or Center of Disease Control (CDC, Atlanta) Adult Mosquito Light Traps. Both types of traps use a normal incandescent light source as an infrared attractant and are programmed to start collections at sunset and terminate at sunrise. Samples are typically retrieved the following morning and forwarded to the laboratory for enumeration and identification. These traps can be augmented (baited) with CO<sub>2</sub>, in canisters, or as dry ice, to increase capture rates as it is another key attractant for female mosquitos. Information gathered from light trap captures can be used to give an indication of the mosquito population size, species complex and the type of development habitat.

New Jersey or CDC Light traps would be deployed to monitor adult mosquito populations in areas with a history of adult mosquito annoyance problems. Benefits associated with these traps include the collection of a much greater number of specimens than with un-baited traps, or from biting counts, and they provide an objective, reproducible sampling method. These collections complement bite count sampling for annoyance by allowing field personnel to more effectively collect and identify mosquito species present in a particular area. Correlation of this data over several years with larval monitoring and adult mosquito biting count data allows for continued, increased forecasting of mosquito populations. Larval and adult mosquitos would be identified according to the taxonomic keys of Darsie and Ward (1981) and Wood, Dang and Ellis (1979), and others as appropriate.

### 3.4.3 Mosquito Development in Saratoga Beach and Black Creek

The largest mosquito development sites at Saratoga Beach and Black Creek are tidally-influenced ponds, channels, ditches and temporarily flooded depressions in salt marsh habitats located in two primary locations: the “Driftwood Marsh”, an old channel/slough extending from Black Creek northwest for some 900 – 1000m, and the Pacific Playgrounds salt marsh habitats at the end of Clarkson Drive south of the Pacific Playgrounds Resort and Marina, ( Figures 1 and 2). The extent and frequency of *Aedes dorsalis* larval development in these habitats occurs in response to a number of influences including precipitation, temperature, tidal fluctuations and resultant seepage. Developing in a synchronous response to fluctuating water levels *Ae. dorsalis* mosquitos hatch throughout the salt marsh and are most numerous during the first half of a season when flooding is also greatest, typically April through July. Although largely confined within a “channel”, direct connection to Georgia Strait is permitted through several culverts and a natural channel at Black Creek. Several days of sustained tides exceeding 4.6 metres (15 ft, measured at Mitlenatch Island), fills the channels and flows through the culverts just north of Black Creek to flow northwest. Resultant flooding, direct and through seepage causes water accumulations in depressions, ponds and channels scattered throughout the marsh. Similar flooding occurs in the Pacific Playgrounds old channel/slough area. Egg eclosion (hatching) occurs within hours of inundation and over the next several days larval development occurs throughout the Driftwood and Pacific Playgrounds salt marshes.



Regular monitoring of salt marsh habitats and correlations with tidal heights during the season ensures that larval mosquito development does not proceed unchecked. With larval populations averaging between 50-100 larvae/350ml dip sample, multiple hatches in a season, and an area amounting to upwards of 95% of the total mosquito development habitat currently identified at Saratoga Beach and Black Creek, the control of larval development in salt marshes is essential to prevent widespread adult mosquito annoyance.

Winter snowpack accumulations and local weather conditions, including total precipitation and frequency, humidity and winds also impact on mosquito development and survival rates. Temperatures have the single greatest impact on the onset and rate of larval development. Depending on local weather conditions temporary and slow-draining, or permanent ponds provide ideal conditions for mosquito development. Producing *Aedes* mosquitos from April through late June, they are a recognizable source of adult mosquito nuisance for area residents. Subsequent larval development, which includes *Culiseta* and *Culex* mosquitos, also causes localized annoyance which may extend through to July and August.



Smaller depressions and non-flowing drainage ditches, tire ruts and artificial containers such as livestock watering-troughs, old tires, landscape, display and irrigation ponds provide the remainder of larval mosquito habitats for the area. Although these sites may often be small, their locations and distribution near outdoor work sites, recreational areas, business' and residential properties makes them important sources of localized adult mosquito annoyance if not effectively monitored and controlled. Adult mosquito specimens collected

during the 2020 Mosquito Development Site Survey suggest that manmade and natural, freshwater development sites locally are capable of producing a variety of species including: *Aedes cinereus*, *Aedes mercurator*, *Aedes provocans*, *Aedes sticticus*, *Culex tarsalis* and *Anopheles punctipennis*.

Although their populations and individual development site sizes are not usually as large as the synchronous hatching *Aedes spp.* mosquitos in salt marsh flood and seepage water habitats, *Culex*, *Culiseta* and *Anopheles* mosquitos make use of a large variety of habitats, including catch basins and containers such as stored tires and equipment, livestock watering troughs, buckets, bird baths and slow flowing streams and ditches. When discovered by field personnel during site inspections these man-made sites would be brought to the attention of property owners for removal, drainage or regular water changes which would eliminate their potential as sources of mosquito development and annoyance.



### 3.5 Mosquito Control Options

Simple but established key elements of a sound IPM approach for this, and every, effective and environmentally compatible nuisance and vector mosquito surveillance and control program are:

- 1) Assessment of need for control using scientifically sound evaluations,
- 2) Development and applications of the best site-specific controls, and;
- 3) As a feedback mechanism, assess the results achieved and adapt as required.

Mosquito development varies from year to year and throughout the season depending on environmental conditions and habitat availability. Environmental cues interact to affect both the timing and magnitude of mosquito development, and adult mosquito survival. These factors include overall development site water levels, fluctuations, water and ambient temperatures, humidity, and precipitation.

Each mosquito development site will have its own unique requirements and treatment options. The PMP for this mosquito control program uses a combination of techniques, and an Integrated Pest Management approach, to achieve the management and control of mosquito populations. The best choice for control reduces both mosquito populations, and the potential for adverse effects on people, domestic animals, livestock and natural ecosystems. Sometimes, particularly with man-made habitats such as ditches, irrigation or display ponds and containers, larval mosquito populations can be reduced or effectively limited using physical or natural controls. These control options are discussed here as they would be considered as a potential solution prior to any larvicide applications.

Many of the possible physical and biological control options suggested below may be supported and possibly implemented by local public works personnel and landowners. Private property owners with mosquito development habitat are best motivated to become involved in their control program through public education initiatives and through consultations with program personnel. Once educated about mosquitos and their habitats, property owners can undertake steps to reduce or eliminate larval mosquito habitat and adult mosquito annoyance on their property. A reduction in larval populations contributes to the overall decrease in adult mosquito annoyance.

The preservation or enhancement of balanced wetland habitats has the best opportunity for a meaningful long-term contribution to overall mosquito control program success through reduction of mosquito populations and enhancement of natural controls including insect, fish and birds. Elimination of stagnant water and enhancements in natural or created ecosystems will be of benefit to overall control program efficacy through decreasing larval habitats and increasing natural mosquito predators. The use of a biorational control products such as *Bacillus thuringiensis* var. *israelensis* (VectoBac 200G / AquaBac 200G) and *Lysinibacillus sphaericus* (VectoLex CG and WSP) when natural control are insufficient, or absent, maximizes the effectiveness and environmental compatibility of the program. IPM-focused mosquito control programs do not have deleterious effects on humans, domestic pets and livestock, wildlife, fish and their food and are routinely conducted throughout British Columbia. There are three larval mosquito control options available to the program. These are physical, biological and biorational product oriented.

### 3.5.1 Physical Source Reduction and Site Modification

A continued focus for the control program technicians and public education initiatives would be the identification, and reduction or elimination, of larval mosquito development habitats wherever possible. Residents and business operators are encouraged to remove, or alter, standing waters which provide suitable habitat for larval mosquito development. For most property owners this involves eliminating water-holding containers, such as buckets and boats or canoes and the draining, or regular changes of water in bird baths, livestock watering troughs, unused wading pools and display ponds.

When done by the homeowners, this permits residents an opportunity to actively participate in their control program. This can be especially important for residents, as two of the most common West Nile virus vector mosquitos, *Culex tarsalis* and *Culex pipiens*, make ready use of manmade habitats, including containers. Installation and maintenance of window screens, Mosquito Magnets (adult mosquito traps) and the use of mosquito repellents by individuals provides additional protection from adult mosquito annoyance and potential disease transmission.



Residents, workers and visitors should minimize outdoor activity at dusk and dawn, wear light-coloured, loose-fitting clothing and minimize the use of fragrant shampoos, perfumes and colognes to further reduce potential adult mosquito nuisance.

Maintaining permanent ditches so they are clear of obstructions or vegetation, replacing failed culverts or grading to effect flow may increase flow, drainage or access by fish or aquatic insect predators. Ditching, grading or filling of roadside depressions may be a suitable solution to decreasing, or preventing, larval development by reducing an area's potential to retain water. When completed as part of routine maintenance activities by public works crews they can be effective means of suppressing local mosquito populations. Any such activities along public roadways, in parks or other publicly owned properties would be coordinated through the appropriate public works and engineering departments. The BCMOE, DFO and other government regulatory agencies, as appropriate, may also need to be consulted prior to any such planned work in area ditches.

Removal or alteration of mosquito producing habitat does not necessarily mean drainage resulting in habitat destruction for other organisms and natural predators such as birds and fish. As part of a comprehensive approach to mosquito control, property owners are encouraged to manage stagnant and non-flowing waters to minimize their use as sources for mosquito development. For example, the removal of emergent shoreline vegetation, combined with either water level management at greater than one metre in depth or a shoreline groomed to a gradient of 3:1 or steeper, effectively eliminates mosquito production in irrigation and settling ponds or other water impoundments. The installation of fountains in man-made golf course and park ponds can reduce their suitability and use as larval mosquito development habitat.



Mosquitos require water to develop, and any efforts to reduce or eliminate standing or stagnant waters, particularly in depressions, tire ruts and containers will prevent larval development and subsequent adult mosquito nuisance. Source reduction around homes and businesses can be easily achieved by residents and owners, allowing them to actively participate in their mosquito control program.

Adult mosquito collection devices such as Mosquito Magnets, which use propane to generate CO<sub>2</sub> will collect adult mosquitos and are marketed by several companies for use by property owners. With a collection range of about ½ hectare (one acre), their ability to reduce mosquito populations sufficiently to provide relief from annoyance on a community level is unlikely without the deployment of numerous units. Their use at a single property/residence though, can have a noticeable impact by collecting adult mosquitos and reducing annoyance.

### 3.5.2 Biological Control

Biological control involves the use of predators, pathogens, and parasites to reduce mosquito populations. Insects predators, both aquatic (i.e. dragon flies, beetles) and terrestrial (i.e. spiders, wasps), contribute to the natural mortalities of both larval and adult mosquitos. Conserving, or enhancing natural habitats wherever possible, allows these predators to contribute to control program effectiveness.



Of all the various predator control methods tested, only larvivorous fish are used operationally in widespread programs. Regan *et al.* (1982) evaluated the effects of three-spined stickleback fish (*Gasterosteus aculeatus*) on mosquito larvae located in the Fraser Valley. They were found to be effective in reducing larval populations. Their natural fecundity combined with their ubiquitous nature makes these fish an ideal natural (biological) control agent. They are a common occurrence in many of ditch systems.

Introduction of fish (Koi, goldfish) to manmade, self-contained outdoor display or irrigation ponds may also reduce, or eliminate larval mosquito development in such habitats. Most practical in the warm, lower mainland Fraser Valley and Vancouver Island, in areas with very cold winters, this type of control requires considerable work and cost which many include the over-wintering of fish indoors or annual replacement. The relocation, or introduction of fish to any natural water course requires approval and permitting through various governmental agencies including Department of Fisheries and Oceans and the BC Ministry of Environment.



Although flying insects can form a large component of the diet for flying insectivores (e.g. bats, swallows, Purple Martins), there is no evidence which suggests they provide a detectable level of mosquito control. Both birds and bats are also opportunistic feeders, and adult mosquitos have been identified as a small component (<2%) of their diet, (Fang 2010 and Gonsalves *et.al.*, 2013). They are not however, scientifically recognized as able to provide any real impact on mosquito populations when used solely as a mosquito population control option.

A one-hectare site, the size of 2 football fields, having a larval population density of just 1 larvae/dip sample, can produce 4,285,714 mosquitos. Reported to eat up to 300 mosquitos a day, a total of some +13,300 birds and/or bats would be required to consume the mosquitos emerging from just one hectare of habitat. Larval populations in much of the program area average between 10-30 larvae/dip sample and can often exceed 100 larvae/dip sample. With between 50 – 100+ hectares of treated habitat, and much of it located within 100-200m of residents and businesses, the sheer potential for adult mosquito populations, likely in the billions, would make a reliance on solely natural controls unlikely to have a noticeable impact on annoyance levels for area residents.



Interested residents would still be encouraged, however, to install bat houses or bird nesting boxes if they wish, since it allows individuals to contribute to a comprehensive, integrated mosquito control program, and in some cases may provide residents with a sense of reduced adult mosquito annoyance.

Pathological agents such as viruses and certain parasites have received much research attention, but none of these are commercially available or approved for use in Canada. The closest form of a naturally occurring control agent which can be readily, easily and safely applied to reduce mosquito populations are products made with the soil bacteria, *Bacillus thuringiensis* var. *israelensis* (*Bti*) and *Lysinibacillus sphaericus* (*Lsph*), which have highly specific insecticidal properties. These larvicides are discussed below.

### 3.5.3 Biorational Control

The control program would use VectoBac 200G and VectoLex CG and WSP larvicide products for larval mosquito control, or AquaBac 200G if required. VectoBac, AquaBac, and VectoLex are the closest form of a natural or biological control agent currently available for routine use in operational mosquito control programs. The use of these products maximizes the environmental compatibility of the annual mosquito control program when used in circumstances where other control options such as physical or natural (biological) control are not practical, they support the principles of an IPM approach to control.

Property owners would be consulted with prior to any larvicide applications and for any recommended physical or biological/natural methods. Product brochures, labels, MSDS sheets and

website addresses would be supplied and reviewed to ensure residents, business, and facility operators understand, are comfortable with, and approve, proposed treatments. In the event that a property owner wishes exclusion from the control program this request would be honoured and noted in the development site database. The product VectoBac 200G contains spores and crystals produced by the bacterium (*Bacillus thuringiensis var. israelensis*, Serotype H-14, Strain AM65-52), and the Product AquaBac 200G contains spores and crystals produced by the bacterium (*Bacillus thuringiensis var. israelensis*, Serotype H-14 Strain, BMP-144). Both are classed as a biorational, rather than conventional, pesticides.

VectoBac and AquaBac act on the larval mosquito stomach and must be eaten to be effective. It is very specific, producing rapid lethal effects (within hours) in larval mosquitos. It has no residual activity, does not bio-accumulate and has no impact on beneficial organisms found in mosquito development habitats. Negative or toxic effects on mammals, birds or other wildlife have not been observed. Formulated as a corn cob granule it requires no mixing and is ready to apply by hand or backpack blower. The granule allows the larvicide to penetrate vegetative covers and reach the water surface where the *Bti* is “released” for consumption by mosquito larvae.



VectoBac 200G Granules

VectoBac and AquaBac are recommended by the manufacturer for use in standing water habitats including temporary and permanent pools in pastures and forested areas, irrigation or roadside ditches, natural marshes or estuarine areas, waters contiguous to fish-bearing waters, catch basins and sewage lagoons.

VectoLex CG and WSP also contain naturally occurring, spore-forming soil bacterium. VectoLex contains spores and crystals produced by *Lysinibacillus sphaericus*. It also is classed as a biorational, rather than conventional, pesticide. VectoLex also acts on the larval mosquito stomach and must be eaten to be effective. VectoLex is very specific and produces lethal effects in a narrow range of mosquito species, including *Aedes vexans* and most *Culex* mosquito species. It does not have any effects on man or animals, fish and other insects which may use these aquatic habitats.

Operationally, the important differences between VectoLex and VectoBac or AquaBac are speed of action and persistence in the larval habitat. Larval mortality can take several days for VectoLex versus several hours with VectoBac/AquaBac. This occurs because *L. sphaericus* is more stable, has a slower settling rate in the water column and the unique ability for its spores to germinate, grow and reproduce in dead mosquito larvae. This is known as recycling and is the mechanism which allows VectoLex to provide long-term, extended control of recurring larval mosquito development. VectoLex CG and WSP are recommended by the manufacturer for use in standing water habitats including temporary and permanent pools in pastures and woodlots, irrigation or roadside ditches, natural marshes or estuarine areas, waters contiguous to fish-bearing waters, catch basins and sewage lagoons.

The use of *Bti* and *Lsph* maximizes the environmental compatibility of the annual mosquito control program since both products are species (target) selective and non-toxic to other aquatic organisms which co-exist in these habitats including insects, fish and amphibians. When used in circumstances where other control options such as physical or cultural control are not practical, they support the principles of an IPM approach to mosquito control.



Extensive product information for VectoBac and VectoLex products can be found at the manufacturer's website [www.valentbiosciences.com](http://www.valentbiosciences.com). The product manufacturer/distributor for AquaBac 200G is AFA Environnement Inc. Additional information on all the larvicide products proposed for this program can be found through the Health Canada, Pest Management Regulatory Agency (PRMA) website [www.pmra-arlc.gc.ca](http://www.pmra-arlc.gc.ca) and the Pesticide Label Search [www.hc-sc.gc.ca](http://www.hc-sc.gc.ca).

### 3.5.4 Chemical Control

Chemical control products and equipment are predominantly used for the purposes of reducing adult mosquito populations. As with most adult insect control programs, adult mosquitos are typically controlled using a broad-spectrum (adulticide) insecticide. Although there are 'natural' adult mosquito control products made from chrysanthemum flower extracts (pyrethrins) and their synthetic equivalents, all adulticides only provide temporary control and are typically broad spectrum, having a deleterious effect on any insect which may come in contact with them. Unless regular and routine treatment of 'problem areas' is completed, uncontrolled adult mosquitos developing in other areas will often move into these treated areas to again cause annoyance.

Typically applied from the ground using cold aerosol sprayers or misters, and much less commonly, from the air using helicopters or fixed-wing aircraft, their mode of action is on the nervous system following contact with the organism and absorption across through the exoskeleton. Because they are applied to the air, and the fact they are non-specific, such applications will not only control adult mosquitos which come in contact with the spray mist, but other non-target organisms such as moths, flies, flying beetles and other insects. Restrictions on applications include habitat type, timing of applications, mosquito population thresholds weather conditions and areas of identified avoidance.

Because of the variable dispersion patterns of mosquitos, geograpy, types of vegetation encountered and ambient weather conditions at the time of treatment, it is difficult to provide any more than temporary control of localized adult mosquito annoyance. Unless regular and routine treatment of 'problem areas' is completed, uncontrolled adult mosquitos developing in other areas will often expand into these treated areas to again cause annoyance.





Adulticide applications **are not** a component of the annual mosquito control program for the CVRD. The mosquito control program described within this PMP does not utilize any chemical control methods for the abatement (control) of larval or adult mosquitos. The Comox Valley Regional District Mosquito Population Management and Control Program would continue to use only the least toxic, most environmentally sound control products available and focus on larval mosquito prevention, reduction and control. As new products become available and registered in Canada, their suitability for use in annual control program will be reviewed.

- **Pesticides and repellents**

Citronella candles, mosquito coils, Konk Automatic Aerosol Sprayers and other such products are marketed as mosquito repellants, or for adult mosquito or biting insect control. These are readily available to residents, campers, and property owners. Property owners may also use items such as these, as required, and as instructed on the product labels.

The mosquito control program would continue to use only the least toxic, most environmentally sound control products available. As new products become available and registered in Canada, their suitability for use in annual control program will be reviewed.



### 3.6 Mosquito Control Program Operations

Approximately 2-4 hectares of potential larval mosquito development habitat have been identified to date ( 2020) in the Saratoga Beach and Black Creek area of the CVRD Electoral Area C. The actual total area that will become infested and require larvicide treatment each season is dependent on hydrological and meteorological events. Fluctuating water levels in floodwater and seepage-influenced development sites, both freshwater and salt marsh (tidal) can cause recurrent larval development with many sites often requiring multiple treatments to effect control. Precipitation, seepage and surface water run-off into permanent or temporary sites can also result in multiple generations of mosquito during a typical season. The well organized, pro-active, integrated pest management approach to mosquito control which has been developed for the District reduces the potential for adult mosquito annoyance by focusing on the identification and timely control of larval populations occurring within the Saratoga Beach and Black Creek region.

In situations where physical alteration of development habitats are impractical or undesirable, or where natural controls are insufficient to reduce nuisance mosquito populations, VectoBac and VectoLex applications would be completed to control larval populations. Control of larvae at their source reduces annoyance levels for area residents and visitors.

As required by the *BC Integrated Pest Management Act and Regulations*, all larvicide applications would be completed, and/or supervised by, personnel certified by BC Ministry of Environment as

pesticide applicators in the category of *Mosquito and Biting Fly Abatement*, or equivalent. All larvicide treatments would be completed using application rates, equipment and methods recommended by the pesticide manufacturer.

### 3.6.1 Public, Worker, and Environmental Safety During Mosquito Control

To ensure public and worker safety, all conditions and restrictions governing biorational larvicide (VectoBac and VectoLex) applications would be followed. Pesticide applicators will follow the guidance of the approved PMP, with regulations contained in the *Pest Control Products Act*, the *Pesticide Control Act*, the *Integrated Pest Management Act and Regulation*, the *Transportation of Dangerous Goods Act* and other relevant government regulations. Larvicide handling, storage and application procedures would conform with those detailed on product labels and endorsed in the '*Pesticide Applicators and Dispensers Handbook*' and associated reference materials supplied through the BC Ministry of Environment.

The "Workplace Hazardous Materials Information System" (WHMIS) is a national system designed to protect Canadian workers from the adverse effects of hazardous materials by providing relevant information. All pest control products intended for use in a workplace require this information on the label or need to have a Safety Data Sheet (SDS) prepared to be eligible for registration. Pesticide handling, storage and application procedures would conform with those detailed on product labels, Material Safety Data Sheets and endorsed in the '*Pesticide Applicators and Dispensers Handbook*' and associated reference materials supplied through the BC Ministry of Environment.

The CVRD mosquito population management and control program is intended to provide residents, workers and visitors relief from extreme or persistent adult mosquito annoyance. The control program is not intended to eliminate the mosquito population and as such landowners and residents who want to be excluded from control operations on their property are recorded and their wishes respected.

Landowner permission to survey, monitor and treat infested larval mosquito habitats located on private property is confirmed each season. Treatment of developing larval mosquito populations in waterbodies on public lands are permitted under this approved PMP. Program personnel will take all practical precautions to protect application personnel, the environment and the general public during all larvicide applications. Prior to any larvicide application field personnel:

- Verify property ownership, treatment site boundaries, public points of access (paths, trails, roadways), pest presence and population size, both pre- and post-treatment.
- Identify AVOID areas, fish-bearing and flowing waters or areas of environmental sensitivity (i.e. bird nesting sites) and the need for, and size of, any Pesticide Free Zones (PFZs) and Pesticide application Buffer Zones (PBZs), if required.
- Community watersheds can be determined by accessing the BC Ministry of Environment Community Watershed listings and informational website:
  - <https://www2.gov.bc.ca/gov/content/environment/air-land-water/water/water-quality/community-watersheds>

- A listing of registered groundwater Wells and Aquifers and an interactive map of active wells is available at:
  - [www.gov.bc.ca/gov/content/environment/air-land-water/water/groundwater-wells-aquifers](http://www.gov.bc.ca/gov/content/environment/air-land-water/water/groundwater-wells-aquifers)
- Larvicides will not be applied to finished drinking water and no Pesticide Free Zones (PFZs) are required for bacterial pesticides as indicated in Section 71(12) of The Integrated Pest Management Act and Regulations. Potable (drinking) water well locations and open water intakes will be identified with the property resident/owner prior to any larvicide treatments.
- Review larvicide product label and comply with recommended precautions for handling and application, safety gear, weather restrictions (wind, temperatures, etc.) and other listed precautions.
- Inform the public of larvicide applications through ongoing public notices, news media articles, advertisements, and personal contact.

### 3.6.2 Larval Mosquito Treatment Thresholds and Application Rates

VectoBac (*Bti*, Strain AM 65-52), AquaBac (*Bti*, Strain BMP-144), and VectoLex (*Lsph*) are only applied when larval mosquitos are present. Larval mosquito surveillance and control protocols would focus efforts on the timely identification and treatment of larval mosquito populations with surveillance and control efforts targeting 1<sup>st</sup> through 3<sup>rd</sup> instar larvae. Given that the most extensive larval development locally involves synchronous hatching *Aedes* mosquitos this strategy ensures maximum control. In addition to treating the most actively growing and feeding instars, it also, allows for retreatment (touch-up) of sites, or portions of sites, that may not have been treated as completely, as desired, because of conditions on the day, changing water levels or because of subsequent hatching. Also, application rates can be lower, and therefore material costs, and overall mosquito larvicide use rates in the environment are reduced. Even though the biorational larvicides containing *Bti* and *Lsph* proposed for use in the program have the safest environmental profiles of any larvicides in common use, decreasing any volume of control product is beneficial and maximizes environmental compatibility.

Treatments targeting mosquito populations with later 3<sup>rd</sup> or 4<sup>th</sup> instars under the guise of allowing natural predators to impact some level of control is not encouraged. It is not recommended on *Bti* or *Lsph* product labels, or by the *Municipal Mosquito Control Guidelines (Ellis, 2005)*. Using biorational larvicides to control mosquito larvae at their source, and as 2<sup>nd</sup> through 3<sup>rd</sup> instar larvae still contributes to the “food web”. Dead larvae become food for many other organisms, including insect detritivores, fungi and bacteria which in turn become food for other aquatic insect and vertebrate predators and grazers. While there may be predation of some mosquito species, such as *Culex spp.* or *Culiseta spp.*, which occur in permanent ponds, and are the two most common larvae in these sites, there are typically zero to very few natural predators in *Aedes* mosquito habitats which are typically dry fields and depressions until inundated with water through snowmelt, precipitation, river seepage, or tidal fluctuations.

Delaying treatments to target mosquito populations with later 3<sup>rd</sup> or 4<sup>th</sup> instars is not ideal. The potential for reduced feeding rates of 4<sup>th</sup> instar larvae may provide incomplete control, reduced efficacy and may result in a number of other undesirable outcomes:

- 1) That larvae develop into the untreatable pupal stage, and then onto nuisance causing adults;
- 2) That field staff may not return at an appropriate time to treat them before this occurs. Changing weather conditions and temperatures over a few days can dramatically accelerate larval development rates; and lastly
- 3) Product manufacturers recommend that later instar larvae are treated with higher application rates, upwards of 10kg/ha (1 gm/metre), thereby requiring more larvicide, increased field personnel time and reduced environmental compatibility.

Changing weather conditions and temperatures over a few days can dramatically accelerate larval development rates; and lastly product manufacturers recommend that later instar larvae are treated with higher application rates, upwards of 10kg/ha (1 gm/M) for VectoBac 200G, and 20 kg/ha for AquaBac 200G (2gm/M), thereby requiring more larvicide, personnel field treatment time, reduced environmental compatibility, and increased cost (\$) per unit treated.

*Aedes* hatch in large numbers, typically +300/dip sample, and inhabit temporary pools created by snowmelt, precipitation, river flood and seepage waters which may only last several days or weeks. These types of temporary habitats seldom have established natural predators and where they may occur, they are typically inadequate to deal with larval populations of such extreme magnitude. A pond the size of a back yard swimming pool (50m<sup>2</sup>), with a larval population of just 1 larvae/350ml dip sample, can produce over 24,000 larvae. A one hectare site, about the size of 2 football fields, with a larval population density of 1 larvae/dip sample can produce 4,285,714 mosquitos.

Typically, upwards of 10 dip samples per development site, depending on site size, are completed. Larger sites will have a greater number of dip samples. Larval mosquito dip samples averaging from 1-3 larvae/350ml dip sample in sites containing predominantly 2<sup>nd</sup> and 3<sup>rd</sup> instar larvae would be the minimum treatment threshold for mosquito larvae found in permanent sites, which typically contain a high proportion of *Culex* and *Culiseta* mosquito larvae. A treatment threshold of five, 1<sup>st</sup> instar larvae/350ml dip sample is utilized when monitoring synchronous, extensive *Aedes spp.* larval development common to early-season, temporary, snowmelt, seepage water and river level influenced habitats. The threshold for 2<sup>nd</sup> and 3<sup>rd</sup> instar *Aedes* larvae would be a minimum of 1 -3 larvae/dip sample. Depending on site location, and its size, and permanence a site may be left untreated if the potential for subsequent adult mosquito nuisance is low, regardless of the larval population size.

VectoLex larvicides, which are largely ineffective against *Aedes* mosquitos would only be utilized to control developing larvae in those permanent and temporary sites having *Culex*, *Culiseta* and *Coquillettidia perturbans* larvae. Larval populations averaging 1-3 larvae/350ml dip sample will be the threshold for treatment using VectoLex larvicides.

These thresholds are based on the “industry standard” used by operational mosquito control programs in the Northwest Mosquito and Vector Control Association (NWMVCA) and American Mosquito Control Association (AMCA). Larval dip sampling, light trap collections and where appropriate, adult mosquito emergence traps would be employed to evaluate post-application larval control results. Larval mortalities of at least 95% would be considered successful. If required, and where indicated by post application sampling, additional, or expanded treatments of nearby areas would be completed to achieve desired efficacy.

VectoBac 200G, AquaBac 200G and VectoLex CG are the larvicides of choice for larvicide applications. The use of *Bti* and *Lsph* maximizes the environmental compatibility of the annual mosquito control program since both products are species (target) selective and non-toxic to other aquatic organisms which co-exist in these habitats including insects, fish and amphibians. When used in circumstances where other control options such as physical or cultural control are not practical, they support the principles of an IPM approach to mosquito control, and they are the most effective, selective, and least persistent larval control agents available.

All Mosquito *Bti* larvicide application rates would be within those recommended by the manufacturer. These rates range from 2.5 to 10.0 kilograms per hectare with applications completed under this PMP to be conducted at rates ranging from 4.0 to 8.5 kilograms per hectare. Typically, VectoBac and VectoLex application rates average 7.5 kg/ha for ground applications. These application rates have been demonstrated as effective under the conditions encountered at Saratoga Beach and Black Creek. Routine application rates for AquaBac 200G are reported to range from 4.0-13.0 kg/ha. Factors influencing application rates include application method, larval age classes (instars) and populations, the density and type of vegetation cover at treatment areas (grasses, deciduous forest), organic matter, water depths etc. All application rates would be within the ranges, and recommendations of the PMRA-approved manufacturers labels. All applications are followed with post-application monitoring to confirm the effectiveness of treatments.

All larvicide applications to small and accessible sites are completed, where required, by hand broadcast or backpack spreader during the mosquito control season. Fluctuating water levels in many of these sites cause repeated larval development requiring repeated treatment. Certified applicators achieve the label recommended application rates (kg/ha) by applying the larvicide granules and at the appropriate concentrations of granules/ft<sup>2</sup>.

Before treating an area, ground applicators review available site maps, estimate the site size (m<sup>2</sup>) and then perform a calculation to determine the volume of VectoBac 200G to be applied. With an application rate of 7.5 kg/ha, and a site size of 1000m<sup>2</sup>, the applicator would measure out 750gms of VectoBac. Applicators then do their best to distribute the granules equally across the water surface while they move around the perimeter of larger and deeper sites, or as they walk through shallower (<30cm deep) sites and as they broadcast the granules by hand or with a back-pack applicator. The desired application rate is achieved by modifying the walking or throttle speed, when using a backpack applicator, or by adjusting the frequency and number of “hand broadcasts” for granules being thrown across the surface by applicators.

For catch basins (CBs), VectoLex WSP 10gm satchels, are applied as one per CB. Alternatively, VectoLex CG can be applied to catch basins. If used, VectoLex CG would be applied using a standard measuring spoon to deliver 10 gm of granules/catch basin. VectoLex WSP used for the treatment of cattail swamps and similar, difficult to access sites, would be applied at a rate of one 10gm satchel for every 5m<sup>2</sup>- 10m<sup>2</sup> surface water. In permanent ponds and stagnant ditches with difficult access because of thick, overgrown, or dense vegetation (i.e. blackberries and Typha sp. cattails), the long-acting VectoLex WSP may be used for treatments. These 10gm satchels (2cm X 2cm) can be readily thrown into these sites where the bio-degradable, glucose-based bag quickly dissolves, and the granules disperse across the water surface. Application to the water surface is at rates of 1 x 10gm satchel/5m<sup>2</sup> to 10m<sup>2</sup> area.



### 3.6.3 Post Application Monitoring

Within 2-96 hours after (post) treatment with VectoBac 200G, larval mortalities would be confirmed through monitoring using a standard 350 ml mosquito dipper. The goal is for larval population reductions of 95%, or to levels with sampling results mostly measured at zero larvae/350ml dip sample and averaging much less than 1 larvae/350ml dip sample. Post-application monitoring confirms treatment success and allows for the 'touch-up' treatment of any areas which may have, for reasons of geography, vegetative cover or access, received inadequate application. Because larval mortality from VectoLex can take several days to occur, and can continue to occur for several weeks, treated larval habitats would be monitored on a regular basis with re-treatment completed as required.

VectoLex post-application monitoring is completed 5-10 days following application, and then weekly to determine if additional treatment is required. VectoLex mortality is not as immediate as VectoBac, and its effectiveness is determined by the absence of developing fourth instar larvae when monitored post application. Because of its recycling in dead mosquito larvae, the appearance of fourth instar larvae indicates that concentration of VectoLex is insufficient to effect control.

Adult mosquito populations would be monitored in harbourage areas adjacent to treated larval development habitats to confirm the effectiveness of larval controls in reducing adult mosquito annoyance. In addition, adult mosquito populations may be monitored at select locations to compare adult mosquito populations between various locations. Given the difference in individual tolerances to mosquito annoyance the success of larval control in limiting adult mosquito populations would be determined through resident reports, interviews and requests for service.

The goal of the annual Mosquito Population and Control Program is to decrease larval mosquito populations sufficiently to reduce, and/or prevent, adult mosquito annoyance for residents, workers and visitors. Property owners and residents are also expected to implement personal protective measures to limit their exposure to adult mosquito annoyance. These include repellent use, clothing choices (long sleeves, light coloured), avoidance of perfumed personal hygiene products (shampoos), window screens and temporal (minimize activity at dusk and dawn) or location avoidance measures which can lessen adult mosquito annoyance.

The general public should be aware that short-lived adult mosquito annoyance may occur during a typical season. The goal of the control program, through pro-active larviciding with VectoBac 200G and VectoLex CG and WSP, is to prevent extreme or persistent, extended adult mosquito annoyance through timely larval surveillance and prevention or control initiatives.

#### 4.0 QUALIFICATIONS OF PROGRAM PERSONNEL

This annual program would be managed by environmental consultants experienced in integrated pest management. The contractor/consultant supplying mosquito control services to the CVRD will have all necessary Pesticide Vendor and Pest Control Service Licences. All personnel working in the annual mosquito control program will be certified pesticide applicators in the category of 'Mosquito and Biting Fly Abatement' or equivalent, as accepted by the BC Ministry of Environment.

Control program management personnel should be Registered Professional Biologists. Field personnel should include University and College graduates or senior Co-Operative Education students studying within the disciplines of biology and environmental science or equivalent practical experience with mosquito population management practices and training.

#### 5.0 LARVICIDE HANDLING AND APPLICATION

As required by the *BC Integrated Pest Management Act and Regulations*, all personnel handling and applying larvicides for the annual mosquito control program must be certified by BC Ministry of Environment as pesticide applicators in the category of *Mosquito and Biting Fly Abatement*, or equivalent. Pesticide applicators will comply with regulations contained within the *Pest Control Products Act*, the *Integrated Pest Management Act*, the *Transportation of Dangerous Goods Act* and other relevant government regulations.

Larvicide handling, storage and application procedures would conform with those detailed on product labels and endorsed in the '*Canadian Pesticide Education Program Applicator Core Manual*', the '*Pesticide Applicators and Dispensers Handbook*' and associated reference materials supplied through the BC Ministry of Environment. This PMP does not attempt to duplicate all the information contained within this handbook and other references. The 'Acts', the Handbook, product labels, manufacturers' websites and any other resource materials detailed in this PMP would be reviewed, as appropriate, before handling, transporting, storing or applying pesticides.

Mosquito control program personnel will take all practical precautions to protect application personnel, the environment and the general public during all pesticide applications. Prior to any pesticide application field personnel are responsible to:

- Review, and as required update, the development site database information for the mosquito control program. The database contains information on property ownership, address, contact telephone number, development site maps, photographs and records of past monitoring and treatment results. The database is continually being updated and

contains information on all known mosquito development habitats, including those located on public and private lands.

- Confirm property ownership, treatment site boundaries, public points of access (paths, trails, roadways), pest presence and population size, both pre- and post-treatment.
- Confirm the status and boundaries of properties where the owner or resident have indicated they wish to be excluded from the mosquito control program (AVOID).
- Confirm the locations of wells or other potable (drinking) water sources. No larvicides will be applied to potable waters for the purposes of mosquito control.
- Inform the public of impending applications through public notices, news media articles, advertisements and/or personal contact with property owners prior proposed treatments.
- Review pesticide product labels and comply with recommended application rates and directions regarding pesticide handling and application including safety gear, weather restrictions (wind, temperatures, etc.) and other listed precautions.

The following sections provide details on procedures and protocols which will protect the public and the environment during larvicide transportation, storage, handling and applications. Only bacterial larvicide products are proposed for use in this mosquito surveillance and control program. No Pesticide Free Zones (PFZs) are required for bacterial pesticides as indicated in Section 71(12) of *The Integrated Pest Management Act and Regulations*.

### **5.1 Larvicide Transportation**

During transportation, all pesticides would be secured to prevent an accidental spillage or theft. Granular VectoBac 200G, AquaBac 200G, and VectoLex larvicide products would be secured and handled to prevent tearing of bags, spillage and exposure to adverse weather conditions such as precipitation. They will be transported in secure lockable compartment such as a vehicle trunk, a locked storage box, container or truck canopy.

Applicators would typically only transport the minimum amounts of pesticide required to complete the proposed treatments. With granular products, it is common for field personnel to require less than forty kilograms of VectoBac 200G, AquaBac 200G, or VectoLex CG for a typical workday. Applicators will carry within their vehicles a suitable spill clean-up kit, basic first aid and appropriate personal protective safety gear and supplies, as required. For granular products, this is typically a broom and dustpan. Any spilled larvicide would be recovered and applied to active larval treatment sites.

Larvicides would not be transported in the passenger compartment of a vehicle and would remain separate from food, clothing or similar items during transport. Any applicator who has product stolen or removed from his/her vehicle would follow the notification procedures for the appropriate authorities immediately, including police.

## 5.2 Larvicide Storage

The CVRD may make available a secure, dry, well-ventilated pesticide storage space for mosquito control larvicides (VectoBac 200G and VectoLex) within their secure public works facilities or equivalent. No large volumes of larvicide are stored on-site over the winter. In an average year, less than 100 kg of VectoBac and VectoLex larvicide may be stored on-site to be available for program start-up in late April.

Emergency telephone numbers for police, fire, ambulance, CANUTEC, Dangerous Goods Emergency Spills, Poison Control, and the BC Ministry of Environment are posted on-site at the storage facility and available at Public Works offices

## 5.3 Larvicide Mixing, Loading and Application

Applicators will follow the directions and precautions warranted by pesticide use as described above and in relevant references. VectoBac and VectoLex granular larvicides are formulated as 'ready to apply'. No mixing is required. They are supplied in thick, plastic bags. All used and empty bags would be disposed of in municipal or regional landfills as directed by the manufacturer on the PMRA approved pesticide label and SDS sheets. Field personnel would wear suitable safety gear, including the appropriate respirator/dust masks, ear protection, rubber gloves, boots and other protective equipment as indicated by larvicide labels, SDS sheets, and manufacturer instructions.

All handling of pesticides would be conducted in level, well ventilated, outside areas under conditions of minimal winds and no precipitation. In the event of accidental spillage personnel would follow accepted spill containment, clean-up and reporting procedures. With granules this typically involves recovery with brooms and dustpans or shovels. This 'recovered' larvicide would be used for the treatment of intended habitats.

Weather forecasts would be consulted, and current weather conditions (wind speed, temperature, precipitation) would be noted, and recorded, during all larvicide applications. In the event that wind speeds during larvicide applications are sufficient to cause the displacement, or drift, of granular larvicides outside of the treatment area, applications would be suspended until suitable conditions return. Similarly, should precipitation be sufficient to cause larvicide (corn cob) granules to clump and clog equipment (backpack blowers), applications will be suspended until suitable conditions return.

Applications of VectoBac 200G and VectoLex to within 10 metres of fish-bearing waters and to waters contiguous to fish-bearing waters is anticipated and as permitted on the Health Canada, Pesticide Regulatory Management Agency (PRMA) approved product labels. Permanent fish-bearing and flowing waters would not be treated. A minimum 10 metre Pesticide Free Zone (PFZ) will be maintained from all potable open water sources, with other buffer zones as required through authorization by the BCMOE. As required by the BC *Integrated Pest Management Act* all larvicide applications are completed by personnel certified by BC Ministry of Environment as pesticide applicators in the category of *Mosquito and Biting Fly Abatement*, or an equivalent, appropriate category.

#### 5.4 Equipment Maintenance and Calibration

Ground-based applications of VectoBac 200G and VectoLex CG are completed by hand broadcast or motorized back-pack type (leaf blower) applicator. Applicators would adjust their walking speed, and throttle speed if using backpack blowers, to ensure they are achieving the correct application rate/density of granules per square foot of water surface. For an application rate of 7.5 kg/ha this is 4-5 granules per square foot.

#### 6.0 CONTROL PROGRAM SYNOPSIS

The mosquito population management and control program developed for the Comox Valley Regional District utilizes a pro-active, integrated approach which focuses on larval mosquito control. The CVRD Mosquito Population Management and Control Program is not intended, nor would it be possible, to eliminate the local mosquito population. An appropriate scope of operations and the prevention or timely treatment of larval mosquitos at their source will reduce local adult mosquito populations.

Public education during the term of this PMP will involve regular news media exposure, public information meetings, pamphlets, doorknob hangers and notice boards, website or social media posts, field personnel interaction with residents, visitors and business operators. These initiatives increase the general public's awareness of program operations and goals and encourages the general public to report adult mosquito annoyance, potential larval development sites and to have input into their control program.


Regular monitoring and treatment of larval mosquitos is a key element to mosquito control program success. These development habitats must be identified and regularly surveyed during a control season to ensure timely detection of larval mosquito development. Surveying, monitoring and control of larval mosquito infestations would begin in April and continue through August, and possibly September, depending on conditions. Program methodologies would continue to concentrate on larval control initiatives with a goal to reduce the extent of standing water development habitats. Developing larval mosquito populations would be controlled through the application of the biorational larvicides VectoBac 200G and VectoLex CG. Adult mosquito population monitoring would be conducted as part of routine control program operations. It allows for the evaluation of larvicide efficacy and provides control personnel with information useful in the location of any previously undetected larval development habitat.

Evaluation of the program in terms of effectiveness and ability to satisfy the needs of the general public is conducted as an on-going process. The cooperation and support of local businesses, ranchers, and property owners is indicative of true community spirit and support for a successful program which benefits workers, residents and visitors to the CVRD Electoral Area C. A well organized, pro-active, integrated pest management approach which concentrates on larval mosquito control ensures a safe, effective and environmentally compatible program.

## 7.0 REFERENCES AND LITERATURE CITED

- American Mosquito Control Association 1973. Elements of Comprehensive Mosquito Control. Mosquito News. Vol. 3, No. 18, 86pp.
- B.C. Centre for Disease Control, March 2004. Arbovirus Surveillance and Response Guidelines for British Columbia. [www.BCCDC.ca](http://www.BCCDC.ca)
- B.C. Centre for Disease Control, 03 July 2008 ,. BC WNv Mosquito Control Working Group excerpt: BC provincial Mosquito Pest Management Plan.
- B.C. Centre for Disease Control website: <http://www.bccdc.ca>
- B.C. Centre for Disease Control, 29 May 2015. Email from Marsha Taylor, Epidemiologist BCCDC.
- B.C. Ministry of Agriculture and Food. Mosquito Control Guide. Queens Printer, Victoria, 1984.
- Belton, Peter. The Mosquitos of British Columbia. Victoria: Handbook (British Columbia Provincial Museum) No 41, 1983, 189 pages.
- Current Environmental. Saratoga Beach and Miracle Beach Nuisance Mosquito Study, March 2017.
- Darsie, R. and Ward, R. , 1981. Identification and Geographical Distribution of the Mosquitoes of North America, North Mexico, American Mosquito Control Association, 313pp.
- Ellis, R.A., 1985. Survey of Approaches Taken to Mosquito Control by other North American Jurisdictions, City of Winnipeg Parks and Recreational Branch, 19pp.
- Ellis, Roy., 21 May 2001. Municipal Mosquito Control Guidelines, Health Canada Bureau of Infectious Diseases, 54 pages.
- Fyfe, M., 2004. Arbovirus Surveillance and Response Guidelines for British Columbia, Draft 4.1., BC Centre for Disease Control, Vancouver, BC, 46p., [www.bccdc.org](http://www.bccdc.org)
- Lacey, L.A., M.S. Mulla., 1988. Safety of *Bacillus thuringiensis sp. israelensis* and *Bacillus sphaericus* to Nontarget Organisms in the Aquatic Environment, 19pp.
- Duka Environmental Services Ltd., 2016 - 2024 District of North Cowichan Nuisance and Vector Mosquito Control Program Summary Report. Supplied Annually.
- Duka Environmental Services Ltd., 2016 - 2024 Central Saanich, Nuisance and Vector Mosquito Control Program Summary Reports. Supplied annually.
- Duka Environmental Services Ltd., 2020 - 2024 Comox Valley Regional District, Nuisance and Vector Mosquito Control Program Summary Reports. Supplied annually.

- Duka Environmental Services Ltd., City of Delta, Pest Management Plan for 2021 - 2026 Nuisance and Vector Mosquito Control Programs. Draft November 2020.
- D.G. Regan and Associates Ltd. (Duka Ltd) , Mosquito Management and West Nile Response Plan based on BC CDC WNV Health Alert, September 2004. Table, 1 page.
- Muhammad Morshed, BCCDC. Sampling/identification of all specimens recommended. Email 13 February 2015
- Mahmut Dogramaci, et al. A method for Subsampling Terrestrial Invertebrates Samples in the Laboratory: Estimating Abundance and Taxa Richness, March 2010
- Rebekah Sudia, Public Health Ecologist, Multnomah County Vector Control District, Portland Oregon. Random sampling and procedures for adult mosquito sample identifications. Email 17 February 2015
- Regan, D.G., et al., 1980. Investigations into the Use of Three Spined Stickleback Fish as a Mosquito Control Agent. Appendix to Lower Mainland Regional Districts' Mosquito Control Board Report, 18pp.
- Regan, D.G., Harvey, D. et al., 1982. Use of *Bacillus var. israelensis* as a mosquito control agent. Can. Journ. Env. Hlth. Review, 14pp.
- Service, R., 1976. Mosquito Ecology: Field Sampling Methods. Halsted Press, Toronto, 583pp.
- Schleier, J.J., 2008. Are Mosquito fish safe? – An ecological Risk Assessment for use in the Northwest., Montana State University, Bozeman, Montana. Presentation 48th Annual Northwest Mosquito and Vector Control Association meeting, Coeur d'Alene,
- Wood, D.M., Dang, P.T., Ellis, R.A., 1979. The Insects and Arachnids of Canada; Part 6, Diptera; Culicidae. Canadian Government Publishing Centre, Ottawa, 390pp



**Figure 1: Saratoga/Miracle Beach; Larval Mosquito Development Site Locations**

# VectoBac® 200G

## BIOLOGICAL LARVICIDE

### GRANULE

GROUP

11

INSECTICIDE

RESTRICTED

**GUARANTEE:**

*Bacillus thuringiensis* subsp. *israelensis*,  
Serotype H-14, strain AM 65-52, 200 International  
Toxic Units (ITU) per milligram (0.2 billion ITU/KG)

REGISTRATION NO. 18158

PEST CONTROL PRODUCTS ACT

List No. 60214-13

**INDEX:**

- 1.0 Precautions
- 2.0 First Aid
- 3.0 Toxicological Information
- 4.0 Storage
- 5.0 Disposal
- 6.0 Notice to User
- 7.0 Directions for Use

**READ THE LABEL BEFORE USING**  
**KEEP OUT OF REACH OF**  
**UNAUTHORIZED PERSONNEL**  
**POTENTIAL SENSITIZER**  
**CAUTION EYE IRRITANT**

**1.0 PRECAUTIONS**

**KEEP OUT OF REACH OF UNAUTHORIZED PERSONNEL**  
**MAY CAUSE SENSITIZATION**  
**CAUTION EYE IRRITANT**

DO NOT apply directly to treated, finished drinking water reservoirs or drinking water receptacles when the water is intended for human consumption.

Avoid contact with skin, eyes, and clothing. Avoid breathing dust/spray mist. Wear a long sleeved shirt, long pants, waterproof gloves, shoes and socks, eye goggles and NIOSH-approved respirator with any N-95, R-95, or P-95 filter for biological products when handling, mixing/loading or applying the product and during all clean-up/repair activities. Applicators may remove gloves, eye goggles and respirators if the design and delivery of the application apparatus reduces exposure to a negligible level (e.g. backpack sprayer with application wands that apply product directly over water surface). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**2.0****FIRST AID**

<b>If on skin or clothing</b>	Rinse skin immediately with plenty of water. Remove contaminated clothing and wash separately before reuse. If irritation occurs and persists or is severe, seek medical attention.
<b>If in eyes</b>	Hold eye open and rinse slowly and gently with water. Remove contact lenses, if present, then continue rinsing eye. If irritation occurs and persists or is severe, seek medical attention.
<b>If inhaled</b>	Move person to fresh air, apply respiration if needed and seek medical attention.
<b>If swallowed</b>	Rinse mouth and throat with copious amounts of water. DO NOT induce vomiting. Promptly contact a physician or poison control centre. DO NOT give anything by mouth to an unconscious person.
<b>General</b>	Seek medical attention if irritation or signs of toxicity occur and persist or is severe. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**3.0****TOXICOLOGICAL INFORMATION**

Treat symptomatically.

**4.0****STORAGE**

In order to ensure microbial purity and potency, VectoBac 200G should be stored in the original container at 0 - 25°C and used within 24 months of the date of manufacture.

**5.0****DISPOSAL**

Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instruction for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**6.0****NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**NATURE OF RESTRICTION:** This product is to be used only in the manner authorized; consult local pesticide regulatory authorities about use permits which may be required.

## 7.0 DIRECTIONS FOR USE

MOSQUITOES	Suggested Range Rate
Habitat: Standing water	3 – 10kg/ha* (0.3 – 1.0 g/m <sup>2</sup> )

Temporary and permanent pools in pastures and woodlots, irrigation or roadside ditches, natural marshes or estuarine areas, water contiguous to fish-bearing water, catch basins and sewage lagoons.

\*Use higher rates in deep and/or polluted water, and when late 3rd and 4th instar larvae predominate.

Apply recommended rate by conventional aerial or ground equipment. Uniform coverage is necessary for best results. For aerial application, apply in uniform non-overlapping swaths when conditions do not favour drift or when wind speeds are less than 10 km/h.

A 3 to 14 day interval between applications should be employed. Monitoring will indicate the appropriate retreatment interval. VectoBac 200G Biological Larvicide does not affect non-target, aquatic, invertebrate predators and parasites which are non-filter feeders. Therefore, longer periods of suppression may result since these beneficials would be conserved to aid in mosquito population management.

**AERIAL APPLICATION INSTRUCTIONS**

Apply only by fixed-wing or rotary aircraft equipment that has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product-specific. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application by using appropriate marking devices and/or electronic guidance equipment.

**Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even coverage.

Apply only when meteorological conditions are in compliance with local and/or provincial authorities.

**Operator Precautions**

DO NOT allow the pilot to mix product to be loaded onto the aircraft. However, loading of premixed product with a closed system is permitted. It is desirable that the pilot has communication capabilities at each treatment site at the time of application. The field crew and the mixer/loaders must wear the personal protective equipment described in the PRECAUTIONS section of this label. When handlers/loaders use closed systems to load product onto the aircraft, the handler requirement for eye goggles and a NIOSH-approved respirator/mask with any N-95, R-95, or P-95 filter for biological products may be waived. When reduced personal protective equipment is worn, the respirator/mask and eye goggles must be immediately available for use in an emergency such as a spill or equipment breakdown. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing must be washed before reuse. Decontaminate aircraft cockpit and vehicle cabs if contamination occurs.

**Product Precautions**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-323-9597 or obtain technical advice from the distributor or from your provincial agricultural or forestry representative. Application of this specific product must meet and/or conform to the aerial uses and rates on this label.

**RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, please note that VectoBac 200G Biological Larvicide contains a Group 11 insecticide. Any insect population may contain individuals naturally resistant to VectoBac 200G Biological Larvicide and other Group 11 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same site. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. The following appropriate resistance management strategies should be followed to delay insecticide resistance:

- Where possible, rotate the use of VectoBac 200G Biological Larvicide or other Group 11 insecticides with different groups that control the same pests in a site.
- Insecticide use should be based on an Integrated Pest Management program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or integrated pest management recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance, contact Valent BioSciences Corporation at 1-800-323-9597.

VectoBac is a registered trademark of Valent BioSciences Corporation, U.S.A.

Registrant:



870 TECHNOLOGY WAY  
LIBERTYVILLE, IL 60048 U.S.A.  
1-800-323-9597

Canadian Agent:  
Valent Canada, Inc.  
6-130 Research Lane  
Guelph, Ontario N1G 5G3 CANADA

04-7021/R2 ©Valent BioSciences Corporation, December 2011

# VectoLex<sup>®</sup> CG

## BIOLOGICAL LARVICIDE

**GRANULAR**

**RESTRICTED**

**GROUP 11A2 INSECTICIDE**

**GUARANTEE:**  
*Bacillus sphaericus* Strain 2362 . . . . . 50 BslTU/mg

REGISTRATION NO. 28008  
 PEST CONTROL PRODUCTS ACT List No. 05722-13

**INDEX:**

- 1.0 Notice to User
- 2.0 Nature of Restriction
- 3.0 Limitations
- 4.0 Restricted Uses: Directions for Use
- 5.0 Mosquito Larval Control
- 6.0 Resistance Management Recommendations
- 7.0 Precautions
- 8.0 First Aid
- 9.0 Toxicological Information
- 10.0 Storage
- 11.0 Disposal
- 12.0 Notice to User

**READ THE LABEL BEFORE USING**  
**PREVENT ACCESS BY UNAUTHORIZED PERSONNEL**  
**POTENTIAL SENSITIZER**  
**CAUTION EYE IRRITANT**

<b>1.0</b>	<p><b>NOTICE TO USER</b></p> <p>This pest control product is to be used only in accordance with the directions on the label. It is an offence under the <i>Pest Control Products Act</i> to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.</p>
<b>2.0</b>	<p><b>NATURE OF RESTRICTION</b></p> <p>This product is to be used only in the manner authorised; consult provincial pesticide regulatory authorities regarding appropriate use permits that may be required.</p>
<b>3.0</b>	<p><b>LIMITATIONS</b></p> <p><b>DO NOT</b> apply directly to treated, finished drinking water reservoirs or drinking water receptacles.</p>
<b>4.0</b>	<p><b>RESTRICTED USES: DIRECTIONS FOR USE</b></p> <p>VectoLex CG is a mosquito larvicide to be applied, without mixing or dilution, by conventional ground or aerial application equipment. Apply to mosquito breeding sites when sampling indicates that mosquito larvae are present. For best results, apply when young larval stages are present. Reapply at a minimum interval of two weeks as needed, if monitoring indicates that further applications are required. Do not reapply within two weeks of application. Do not apply more than six times per treatment site per season.</p> <p><b>Aerial Application Instructions:</b>                  Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the</p>

5.0

atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application by employing appropriate marking devices and/or electronic tracking equipment.

**Use Precautions:**  
 Apply only when meteorological conditions at the treatment site allow for complete and even coverage. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

**Operator Precautions:**  
 Do not allow the pilot to mix product to be loaded onto the aircraft. Loading of premixed product with a closed system is permitted. It is desirable that the pilot has communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear the personal protective equipment described in the PRECAUTIONS section of this label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Product Specific Precautions:**  
 Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-323-9597 or obtain technical advice from the distributor or from your provincial agricultural or forestry representative. Application of this specific product must meet and/or conform to the aerial uses and rates on this label. Rinse and flush spray equipment thoroughly following each use.

**5.0 MOSQUITO LARVAL CONTROL**

VectoLex CG is a mosquito larvicide. It is not effective against mosquito adults and pupae. Apply to mosquito larval breeding sites when sampling indicates that mosquito larvae are present. For best results, apply when young larval stages are present.

For use in:	Mosquito species controlled	Application rate and interval	Application methods
Water bodies: freshwater marshes, salt marshes, flood plains, flooded fields and pastures, wetlands, ponds, storm water detention/retention and seepage ponds, wastewater sewage effluent, sewage lagoons, oxidation ponds, log ponds, impounded waste water, septic ditches, drainage ditches including open storm sewers and irrigation ditches	<i>Culex</i> spp. <i>Culiseta</i> spp. <i>Aedes vexans</i> (Other <i>Aedes</i> spp. and <i>Ochlerotatus</i> spp. have variable degrees of susceptibility to VectoLex CG)	5.6-16.8 kg product/ha (0.56-1.68 g product/m <sup>2</sup> ) of water surface area.  Use the higher rate in water polluted with sewage, water with high organic content and water with a high level of suspended solids.  Reapply at a minimum interval of two weeks as needed, if monitoring indicates that further applications are required. Do not reapply within two weeks of application. Do not apply more than six times per treatment site per season.	Apply by ground or aerial application equipment capable of uniform delivery of VectoLex CG Biological Larvicide over the water surface.
Waste tires	<i>Culex</i> spp. <i>Culiseta</i> spp. <i>Aedes triseriatus</i>	0.56-1.68 g product/m <sup>2</sup> of water surface area. Use the higher rate in water with high organic content and water with a high level of suspended solids.	Apply by hand or ground application equipment to individual tires which contain standing water.

## 5.0 MOSQUITO LARVAL CONTROL (CONT'D)

For use in:	Mosquito species controlled	Application rate and interval	Application methods
Waste tires	<i>Culex</i> spp. <i>Culiseta</i> spp. <i>Aedes triseriatus</i>	Do not reapply within one week of application. Reapply at a minimum interval of one week as needed, if monitoring indicates that further applications are required.	Use with other mosquito management techniques such as shredding waste tires, removing standing water and covering the stacked tires.
Water bodies: freshwater marshes, flood plains, flooded fields and pastures, wetlands, ponds, storm water detention/retention and seepage ponds, wastewater sewage effluent, sewage lagoons, oxidation ponds, log ponds, impounded waste water, septic ditches, drainage ditches including open storm sewers and irrigation ditches	<i>Coquillettia perturbans</i> (may reduce the number of larvae)	8-16.8 kg product /ha (0.8-1.68 g product/m <sup>2</sup> ) of water surface area.  Use the higher rate in water polluted with sewage, water with high organic content and water with a high level of suspended solids.  Reapply at a minimum interval of two weeks after application; however, the effects on larval numbers may not be evident for up to three weeks after application.  Reapplication should be based on larval monitoring. Do not reapply within two weeks of application. Do not apply more than six times per treatment site per season.	Apply by ground or aerial application equipment capable of uniform delivery of VectoLex CG Biological Larvicide over the water surface.

## 6.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

Mosquito populations may contain individuals naturally resistant to VectoLex CG. The resistant individuals may dominate the mosquito population if VectoLex CG is used repeatedly as the sole means of control in the same geographic location/use area. To delay/avoid the resistance of mosquito populations to VectoLex CG it is recommended that users:

- Rotate the use of VectoLex CG with other mosquito larvicides currently registered in Canada, which do not contain *Bacillus sphaericus* as the active ingredient, providing they are registered for use in control of the same pests in the same sites.
- Treat a portion of the target area with a *Bti* formulation or an alternative insecticide ensuring the continual existence of populations of mosquitoes not exposed to VectoLex CG within a given geographic location.
- Insecticide use in mosquito control should be based on an IPM program that includes scouting, record keeping, and considers cultural/habitat, biological and chemical control practices suitable for the area to be treated.
- Monitor treated pest populations for resistance development.
- For further information or to report suspected resistance contact Valent BioSciences Corporation at 1-800-323-9597 or at [www.valentbiosciences.com](http://www.valentbiosciences.com).

Registrant:



870 TECHNOLOGY WAY  
LIBERTYVILLE, IL 60048 U.S.A.  
1-800-323-9597

Canadian Agent:  
Valent Canada, Inc.  
6-130 Research Lane  
Guelph, Ontario N1G 5G3 CANADA

## 7.0 PRECAUTIONS

## PREVENT ACCESS BY UNAUTHORIZED PERSONNEL.

May cause sensitization. May irritate eyes. Avoid contact with skin, eyes or clothing. Mixer/loaders and applicators not in enclosed cabs or aircraft must wear a long-sleeved shirt, long pants, shoes plus socks, eye goggles, waterproof gloves and a dust/mist filtering respirator (MSH/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N-95, R-95, P-95 or HE filter for biological products when handling, mixing/loading or applying the product and during all cleanup/repair activities. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

## 8.0 FIRST AID

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**GENERAL: IMMEDIATELY** seek medical attention if irritation or signs of toxicity occur and persist or are severe.

## 9.0 TOXICOLOGICAL INFORMATION:

Treat symptomatically.

## 10.0 STORAGE

Store at temperatures between 0°C and 25°C. Store container upright and keep tightly closed when not in use. Material must be used within 12 months of the Date of Manufacture.

## 11.0 DISPOSAL

1. Completely empty the bag into the application equipment.
  2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
  3. Make the empty bag unsuitable for further use.
  4. Dispose of the bag in accordance with provincial requirements.
- For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.
- DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

## 12.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

*VectoLex* is a registered trademark of Valent BioSciences Corporation, U.S.A.